

## CO027 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10811b

# **Specification**

# CO027 Antibody (C-term) - Product Information

Application WB, FC,E
Primary Accession Q2M3C6

Other Accession Q9N0B5, NP\_689548.2

Reactivity Human, Mouse

Predicted Monkey
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 58444
Antigen Region 503-531

# CO027 Antibody (C-term) - Additional Information

#### Gene ID 123591

### **Other Names**

Transmembrane protein C15orf27, C15orf27

## Target/Specificity

This CO027 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 503-531 amino acids from the C-terminal region of human CO027.

# **Dilution**

WB~~1:1000 FC~~1:10~50

#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

CO027 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# CO027 Antibody (C-term) - Protein Information

Name TMEM266 (<u>HGNC:26763</u>)





**Function** Voltage-sensor protein present on the post-synaptic side of glutamatergic mossy fibers and granule cells in the cerebellum (PubMed:<u>25165868</u>, PubMed:<u>30810529</u>). Despite the presence of a voltage- sensor segment, does not form a functional ion channel and its precise role remains unclear (PubMed:<u>25165868</u>, PubMed:<u>30810529</u>). Undergoes both rapid and slow structural rearrangements in response to changes in voltage (PubMed:<u>30810529</u>). Contains a zinc-binding site that can regulate the slow conformational transition (PubMed:<u>30810529</u>).

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein. Cell projection, dendrite Perikaryon. Note=Present in the dendrites and soma of cerebellar granule neurons, but not in their axon

### **Tissue Location**


Mainly expressed in the cerebellum (PubMed:25165868). Also expressed in cerebral cortex, skeletal muscle and thyroid, but at much lower levels (PubMed:25165868)

# CO027 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# CO027 Antibody (C-term) - Images




CO027 Antibody (C-term) (Cat. #AP10811b) western blot analysis in MCF-7 cell line lysates (35ug/lane). This demonstrates the CO027 antibody detected the CO027 protein (arrow).





CO027 Antibody (C-term) (Cat. #AP10811b) western blot analysis in mouse Neuro-2a cell line lysates (35ug/lane). This demonstrates the CO027 antibody detected the CO027 protein (arrow).



CO027 Antibody (C-term) (Cat. #AP10811b) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

# CO027 Antibody (C-term) - References

Zody, M.C., et al. Nature 440(7084):671-675(2006)