

DNAJC22 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10871B**Specification**

DNAJC22 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q8N4W6
Other Accession	Q17QW0 , NP_079178.2
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	38086
Antigen Region	247-276

DNAJC22 Antibody (C-term) - Additional Information**Gene ID** 79962**Other Names**

DnaJ homolog subfamily C member 22, DNAJC22

Target/Specificity

This DNAJC22 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 247-276 amino acids from the C-terminal region of human DNAJC22.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DNAJC22 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DNAJC22 Antibody (C-term) - Protein Information**Name** DNAJC22

Function May function as a co-chaperone.

Cellular Location


Membrane; Multi-pass membrane protein

DNAJC22 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DNAJC22 Antibody (C-term) - Images



DNAJC22 Antibody (C-term) (Cat. #AP10871b) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the DNAJC22 antibody detected the DNAJC22 protein (arrow).



DNAJC22 antibody (C-term) (Cat. #AP10871b) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the DNAJC22 antibody (C-term) for

immunohistochemistry. Clinical relevance has not been evaluated.

DNAJC22 Antibody (C-term) - Background

May function as a co-chaperone.

DNAJC22 Antibody (C-term) - References

Behr, M., et al. Nat. Cell Biol. 9(7):847-853(2007)