

**PMAT(Slc29a4) Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP1087b**

**Specification**

---

**PMAT(Slc29a4) Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q8R139</a>
Other Accession	<a href="#">Q7RTT9</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	452-481

**PMAT(Slc29a4) Antibody (C-term) - Additional Information**

**Gene ID** 243328

**Other Names**

Equilibrative nucleoside transporter 4, Solute carrier family 29 member 4, Slc29a4, Ent4

**Target/Specificity**

This PMAT(Slc29a4) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 452-481 amino acids of mouse PMAT(Slc29a4).

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PMAT(Slc29a4) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**PMAT(Slc29a4) Antibody (C-term) - Protein Information**

**Name** Slc29a4 {ECO:0000312|MGI:MGI:2385330}

**Function** Electrogenic voltage-dependent transporter that mediates the transport of a variety of endogenous bioactive amines, cationic xenobiotics and drugs (PubMed:[16873718](#), PubMed:[23255610](#)). Utilizes the physiologic inside-negative membrane potential as a driving force

to facilitate cellular uptake of organic cations (By similarity). Functions as a Na(+)- and Cl(-)-independent bidirectional transporter (By similarity). Substrate transport is pH-dependent and enhanced under acidic condition, which is most likely the result of allosteric changes in the transporter structure (PubMed:[16873718](#)). Implicated in monoamine neurotransmitters uptake such as serotonin, dopamine, adrenaline/epinephrine, noradrenaline/norepinephrine, histamine and tyramine, thereby supporting a role in homeostatic regulation of aminergic neurotransmission in the central nervous system (PubMed:[23255610](#)). Also responsible for the uptake of bioactive amines and drugs through the blood-cerebrospinal fluid (CSF) barrier, from the CSF into choroid plexus epithelial cells, thereby playing a significant role in the clearance of cationic neurotoxins, xenobiotics and metabolic waste in the brain (PubMed:[23255610](#)). Involved in bidirectional transport of the purine nucleoside adenosine and plays a role in the regulation of extracellular adenosine concentrations in cardiac tissues, in particular during ischemia (PubMed:[16873718](#)). May be involved in organic cation uptake from the tubular lumen into renal tubular cells, thereby contributing to organic cation reabsorption in the kidney (PubMed:[23255610](#)). Also transports adenine and guanidine (PubMed:[16873718](#)).

#### **Cellular Location**

Cell membrane {ECO:0000250|UniProtKB:Q7RTT9}; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Note=Localized to the apical blood-cerebrospinal fluid(CSF)-facing membrane of the choroid plexus epithelium

#### **Tissue Location**

Expressed in heart (PubMed:16873718). Expressed in choroid plexus (PubMed:23255610).


#### **PMAT(Slc29a4) Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **PMAT(Slc29a4) Antibody (C-term) - Images**





Western blot analysis of PMAT(Slc29a4) (arrow) using rabbit polyclonal PMAT(Slc29a4) Antibody (C-term) (Cat.#AP1087b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the PMAT(Slc29a4) gene (Lane 2) (Origene Technologies).

#### **PMAT(Slc29a4) Antibody (C-term) - Background**

PMAT(Slc29a4) is a member of the SLC29 family and encodes a plasma membrane protein with 11 transmembrane helices. This protein catalyzes the re-uptake of monoamines into presynaptic neurons, thus determining the intensity and duration of monoamine neural signaling. It has been shown to transport several compounds, including serotonin, dopamine, and the neurotoxin 1-methyl-4-phenylpyridinium.

#### **PMAT(Slc29a4) Antibody (C-term) - References**

Barnes,K., Circ. Res. 99 (5), 510-519 (2006)  
Engel,K., J. Biol. Chem. 279 (48), 50042-50049 (2004)