

SLFN12L Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10917a**Specification**

SLFN12L Antibody (N-term) - Product Information

Application	WB, FC,E
Primary Accession	Q6IEE8
Other Accession	NP_001138499.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	67096
Antigen Region	95-123

SLFN12L Antibody (N-term) - Additional Information**Gene ID** 100506736**Other Names**

Schlafen family member 12-like, SLFN12L, SLFN5

Target/Specificity

This SLFN12L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 95-123 amino acids from the N-terminal region of human SLFN12L.

Dilution

WB~~1:1000

FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SLFN12L Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SLFN12L Antibody (N-term) - Protein Information**Name** SLFN12L**Synonyms** SLFN5


Cellular Location

Membrane; Single-pass membrane protein.


SLFN12L Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SLFN12L Antibody (N-term) - Images

SLFN12L Antibody (N-term) (Cat. #AP10917a) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the SLFN12L antibody detected the SLFN12L protein (arrow).



SLFN12L Antibody (N-term) (Cat. #AP10917a) flow cytometric analysis of Jurkat cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.