

CCNG1 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10939a**Specification**

CCNG1 Antibody (N-term) - Product Information

Application	WB, FC,E
Primary Accession	P51959
Other Accession	Q52QT8 , Q5E9I1 , NP_004051.1 , NP_954854.1
Reactivity	Human
Predicted	Bovine, Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	34074
Antigen Region	20-53

CCNG1 Antibody (N-term) - Additional Information**Gene ID** 900**Other Names**

Cyclin-G1, Cyclin-G, CCNG1, CCNG, CYCG1

Target/Specificity

This CCNG1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 20-53 amino acids from the N-terminal region of human CCNG1.

Dilution

WB~~1:1000

FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CCNG1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CCNG1 Antibody (N-term) - Protein Information**Name** CCNG1

Synonyms CCNG, CYCG1

Function May play a role in growth regulation. Is associated with G2/M phase arrest in response to DNA damage. May be an intermediate by which p53 mediates its role as an inhibitor of cellular proliferation (By similarity).

Cellular Location

Nucleus. Note=DNA replication foci after DNA damage


Tissue Location

High levels in skeletal muscle, ovary, kidney and colon


CCNG1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCNG1 Antibody (N-term) - Images

CCNG1 Antibody (N-term) (Cat. #AP10939a) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the CCNG1 antibody detected the CCNG1 protein (arrow).



CCNG1 Antibody (N-term) (Cat. #AP10939a) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

CCNG1 Antibody (N-term) - Background

The eukaryotic cell cycle is governed by cyclin-dependent protein kinases (CDKs) whose activities are regulated by cyclins and CDK inhibitors. The protein encoded by this gene is a member of the cyclin family and contains the cyclin box. The encoded protein lacks the protein destabilizing (PEST) sequence that is present in other family members. Transcriptional activation of this gene can be induced by tumor protein p53. Two transcript variants encoding the same protein have been identified for this gene. [provided by RefSeq].

CCNG1 Antibody (N-term) - References

Cunningham, J.M., et al. Br. J. Cancer 101(8):1461-1468(2009)
Fornari, F., et al. Cancer Res. 69(14):5761-5767(2009)
Li, H., et al. Mol. Cell. Biol. 29(3):919-928(2009)
Piscopo, D.M., et al. Cancer Res. 68(14):5581-5590(2008)
Seo, H.R., et al. J. Biol. Chem. 283(23):15601-15610(2008)