

**ZCCHC17 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP11017C****Specification**

---

**ZCCHC17 Antibody (Center) - Product Information**

|                   |                             |
|-------------------|-----------------------------|
| Application       | WB, FC,E                    |
| Primary Accession | <a href="#">Q9NP64</a>      |
| Other Accession   | <a href="#">NP_057589.2</a> |
| Reactivity        | Human, Mouse                |
| Host              | Rabbit                      |
| Clonality         | Polyclonal                  |
| Isotype           | Rabbit IgG                  |
| Calculated MW     | 27570                       |
| Antigen Region    | 159-187                     |

**ZCCHC17 Antibody (Center) - Additional Information****Gene ID** 51538**Other Names**

Nucleolar protein of 40 kDa, pNO40, Pnn-interacting nucleolar protein, Putative S1 RNA-binding domain protein, PS1D protein, Zinc finger CCHC domain-containing protein 17, ZCCHC17, PS1D

**Target/Specificity**

This ZCCHC17 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 159-187 amino acids from the Central region of human ZCCHC17.

**Dilution**

WB~~1:1000  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ZCCHC17 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**ZCCHC17 Antibody (Center) - Protein Information****Name** ZCCHC17


**Synonyms** PS1D**Cellular Location**

Nucleus, nucleolus.


**ZCCHC17 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**ZCCHC17 Antibody (Center) - Images**

All lanes : Anti-ZCCHC17 Antibody (Center) at 1:1000 dilution Lane 1: PC-3 whole cell lysate Lane 2: U-2OS whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 28 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



ZCCHC17 Antibody (Center) (Cat. #AP11017c) western blot analysis in mouse NIH-3T3 cell line lysates (35ug/lane). This demonstrates the ZCCHC17 antibody detected the ZCCHC17 protein (arrow).



ZCCHC17 Antibody (Center) (Cat. #AP11017c) flow cytometric analysis of A375 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### **ZCCHC17 Antibody (Center) - References**

Chang, W.L., et al. Biochem. Biophys. Res. Commun. 307(3):569-577(2003)  
Gueydan, C., et al. J. Biol. Chem. 277(47):45034-45040(2002)