

HDAC2 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1102b**Specification**

HDAC2 Antibody (C-term) - Product Information

Application	IF, WB, IHC-P,E
Primary Accession	Q92769
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	456-488

HDAC2 Antibody (C-term) - Additional Information**Gene ID** 3066**Other Names**

Histone deacetylase 2, HD2, HDAC2

Target/Specificity

This HDAC2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 456-488 amino acids from the C-terminal region of human HDAC2.

Dilution

IF~~1:10~50

WB~~1:1000

IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HDAC2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

HDAC2 Antibody (C-term) - Protein Information**Name** HDAC2 {ECO:0000303|PubMed:10545197, ECO:0000312|HGNC:HGNC:4853}**Function** Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) (PubMed:[28497810](#)). Histone

deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (By similarity). Histone deacetylases act via the formation of large multiprotein complexes (By similarity). Forms transcriptional repressor complexes by associating with MAD, SIN3, YY1 and N-COR (PubMed:[12724404](#)). Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development (By similarity). Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin (PubMed:[16428440](#), PubMed:[28977666](#)). Component of the SIN3B complex that represses transcription and counteracts the histone acetyltransferase activity of EP300 through the recognition H3K27ac marks by PHF12 and the activity of the histone deacetylase HDAC2 (PubMed:[37137925](#)). Also deacetylates non-histone targets: deacetylates TSHZ3, thereby regulating its transcriptional repressor activity (PubMed:[19343227](#)). May be involved in the transcriptional repression of circadian target genes, such as PER1, mediated by CRY1 through histone deacetylation (By similarity). Involved in MTA1-mediated transcriptional corepression of TFF1 and CDKN1A (PubMed:[21965678](#)). In addition to protein deacetylase activity, also acts as a protein-lysine deacylase by recognizing other acyl groups: catalyzes removal of (2E)-butenoyl (crotonyl) and 2- hydroxyisobutanoyl (2-hydroxyisobutyryl) acyl groups from lysine residues, leading to protein decrotonylation and de-2- hydroxyisobutyrylation, respectively (PubMed:[28497810](#), PubMed:[29192674](#)).

Cellular Location

Nucleus. Cytoplasm


Tissue Location


Widely expressed; lower levels in brain and lung.

HDAC2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


HDAC2 Antibody (C-term) - Images




Confocal immunofluorescent analysis of HDAC2 Antibody (C-term)(Cat#AP1102b) with 293 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red).




Fluorescent confocal image of Hela cell stained with HDAC2 Antibody (C-term)(Cat#AP1102b). Hela cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with HDAC2 primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min). hHDAC2 immunoreactivity is localized to Nucleus significantly.




Western blot analysis of HDAC2 (arrow) using rabbit polyclonal HDAC2 Antibody (C-term) (Cat.#AP1102b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the HDAC2 gene (Lane 2) (Origene Technologies).



Western blot analysis of anti-HDAC2 Antibody (C-term) (Cat.#AP1102b) in 293 cell line lysates (35ug/lane). HDAC2 (arrow) was detected using the purified Pab.



Western blot analysis of anti-HDAC2 Antibody (C-term) (Cat.#AP1102b) in mouse spleen tissue lysates (35ug/lane). HDAC2 (arrow) was detected using the purified Pab.



HDAC2 Antibody (C-term) (Cat. #AP1102b) immunohistochemistry analysis in formalin fixed and paraffin embedded human breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of HDAC2 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

HDAC2 Antibody (C-term) - Background

Histone deacetylase 2 (HDAC2), or transcriptional regulator homolog RPD3 L1, is highly homologous to the yeast transcription factor RPD3 (reduced potassium dependency 3) gene. As in yeast, human HDA2 is likely to be involved in regulating chromatin structure during transcription. It has been implicated to associate with YY1, a mammalian zinc-finger transcription factor, which negatively regulates transcription by tethering RPD3 to DNA as a cofactor. This process is highly conserved from yeast to human.

HDAC2 Antibody (C-term) - References

Choi, Y.B., et al., J. Biol. Chem. 279(49):50930-50941 (2004).
Zhu, P., et al., Cancer Cell 5(5):455-463 (2004).
Longworth, M.S., et al., J. Virol. 78(7):3533-3541 (2004).
Lu, Y., et al., J. Biol. Chem. 278(48):47792-47802 (2003).
Verdin, E., et al., Trends Genet. 19(5):286-293 (2003).