

OR1J4 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11063b**Specification**

OR1J4 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q8NGS1
Other Accession	Q8VGI1 , NP_001004452.1
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	34959
Antigen Region	275-303

OR1J4 Antibody (C-term) - Additional Information**Gene ID** 26219**Other Names**

Olfactory receptor 1J4, HTPCRX01, Olfactory receptor OR9-21, OR1J4

Target/Specificity

This OR1J4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 275-303 amino acids from the C-terminal region of human OR1J4.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR1J4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR1J4 Antibody (C-term) - Protein Information**Name** OR1J4

Function Odorant receptor.

Cellular Location


Cell membrane; Multi-pass membrane protein.

OR1J4 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR1J4 Antibody (C-term) - Images



All lanes : Anti-OR1J4 Antibody (C-term) at 1:1000 dilution Lane 1: HL-60 whole cell lysate Lane 2: MDA-MB-453 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 35 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



OR1J4 antibody (C-term) (Cat. #AP11063b) immunohistochemistry analysis in formalin fixed and paraffin embedded human testis carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the OR1J4 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

OR1J4 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR1J4 Antibody (C-term) - References

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)
Fuchs, T., et al. Genomics 80(3):295-302(2002)
Parmentier, M., et al. Nature 355(6359):453-455(1992)