

OR10Z1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP11424b

Specification

OR10Z1 Antibody (C-term) - Product Information

Application WB,E
Primary Accession Q8NGY1

Other Accession NP_001004478.1

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Rabbit
Polyclonal
Rabbit IgG
284-313

OR10Z1 Antibody (C-term) - Additional Information

Gene ID 128368

Other Names

Olfactory receptor 10Z1, Olfactory receptor OR1-15, OR10Z1

Target/Specificity

This OR10Z1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 284-313 amino acids from the C-terminal region of human OR10Z1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR10Z1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR10Z1 Antibody (C-term) - Protein Information

Name OR10Z1

Function Odorant receptor.



Cellular Location


Cell membrane; Multi-pass membrane protein.

OR10Z1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

OR10Z1 Antibody (C-term) - Images



OR10Z1 Antibody (C-term) (Cat. #AP11424b) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the OR10Z1 antibody detected the OR10Z1 protein (arrow).

OR10Z1 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR10Z1 Antibody (C-term) - References

Shimada, M., et al. Hum. Genet. 128(4):433-441(2010) Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)