

KNG1 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11683a**Specification**

KNG1 Antibody (N-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P01042
Other Accession	NP_001095886.1 , NP_000884.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	71957
Antigen Region	138-166

KNG1 Antibody (N-term) - Additional Information**Gene ID** 3827**Other Names**

Kininogen-1, Alpha-2-thiol proteinase inhibitor, Fitzgerald factor, High molecular weight kininogen, HMWK, Williams-Fitzgerald-Flaujeac factor, Kininogen-1 heavy chain, T-kinin, Ile-Ser-Bradykinin, Bradykinin, Kallidin I, Lysyl-bradykinin, Kallidin II, Kininogen-1 light chain, Low molecular weight growth-promoting factor, KNG1, BDK, KNG

Target/Specificity

This KNG1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 138-166 amino acids from the N-terminal region of human KNG1.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KNG1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

KNG1 Antibody (N-term) - Protein Information

Name KNG1**Synonyms** BDK, KNG

Function Kininogens are inhibitors of thiol proteases. HMW-kininogen plays an important role in blood coagulation by helping to position optimally prekallikrein and factor XI next to factor XII; HMW-kininogen inhibits the thrombin- and plasmin-induced aggregation of thrombocytes. LMW-kininogen inhibits the aggregation of thrombocytes. LMW-kininogen is in contrast to HMW-kininogen not involved in blood clotting.

Cellular Location

Secreted, extracellular space.


Tissue Location

Secreted in plasma. T-kinin is detected in malignant ovarian, colon and breast carcinomas, but not in benign tumors.


KNG1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KNG1 Antibody (N-term) - Images


Western blot analysis of KNG1 (arrow) using rabbit polyclonal KNG1 Antibody (N-term) (Cat. #AP11683a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the KNG1 gene.



Anti-KNG1 Antibody (N-term) at 1:1000 dilution + Ramos whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 72 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



KNG1 Antibody (N-term) (Cat. #AP11683a) immunohistochemistry analysis in formalin fixed and paraffin embedded human breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of KNG1 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



KNG1 Antibody (N-term) (Cat. #AP11683a) flow cytometric analysis of Ramos cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit

secondary antibodies were used for the analysis.

KNG1 Antibody (N-term) - Background

This gene uses alternative splicing to generate two different proteins- high molecular weight kininogen (HMWK) and low molecular weight kininogen (LMWK). HMWK is essential for blood coagulation and assembly of the kallikrein-kinin system. Also, bradykinin, a peptide causing numerous physiological effects, is released from HMWK. In contrast to HMWK, LMWK is not involved in blood coagulation. Three transcript variants encoding different isoforms have been found for this gene.

KNG1 Antibody (N-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Houlihan, L.M., et al. Am. J. Hum. Genet. 86(4):626-631(2010)
Khan, M.M., et al. Am. J. Physiol. Heart Circ. Physiol. 298 (2), H652-H658 (2010) :
Bellucci, F., et al. Br. J. Pharmacol. 158(8):1996-2004(2009)