

TRIM56 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11813b**Specification**

TRIM56 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O9BRZ2
Other Accession	NP_112223.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	81488
Antigen Region	727-755

TRIM56 Antibody (C-term) - Additional Information**Gene ID** 81844**Other Names**

E3 ubiquitin-protein ligase TRIM56, 632-, RING finger protein 109, Tripartite motif-containing protein 56, TRIM56, RNF109

Target/Specificity

This TRIM56 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 727-755 amino acids from the C-terminal region of human TRIM56.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TRIM56 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TRIM56 Antibody (C-term) - Protein Information**Name** TRIM56 {ECO:0000303|PubMed:21289118, ECO:0000312|HGNC:HGNC:19028}**Function** E3 ubiquitin-protein ligase that plays a key role in innate antiviral immunity by

mediating ubiquitination of CGAS and STING1 (PubMed:[21289118](#), PubMed:[29426904](#)). In response to pathogen- and host- derived double-stranded DNA (dsDNA), targets STING1 to 'Lys-63'-linked ubiquitination, thereby promoting its homodimerization, a step required for the production of type I interferon IFN-beta (By similarity). Also mediate monoubiquitination of CGAS, thereby promoting CGAS oligomerization and subsequent activation (PubMed:[29426904](#)). Promotes also TNFalpha-induced NF-kappa-B signaling by mediating 'Lys-63'-linked ubiquitination TAK1, leading to enhanced interaction between TAK1 and CHUK/IKKalpha (PubMed:[35952808](#)). Independently of its E3 ubiquitin ligase activity, positive regulator of TLR3 signaling. Potentiates extracellular double stranded RNA (dsRNA)-induced expression of IFNB1 and interferon-stimulated genes ISG15, IFIT1/ISG56, CXCL10, OASL and CCL5/RANTES (PubMed:[22948160](#)). Promotes establishment of an antiviral state by TLR3 ligand and TLR3-mediated chemokine induction following infection by hepatitis C virus (PubMed:[22948160](#)). Acts as a restriction factor of Zika virus through direct interaction with the viral RNA via its C-terminal region (PubMed:[31251739](#)).

Cellular Location

Cytoplasm.


Tissue Location

Widely expressed (at protein level).

TRIM56 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRIM56 Antibody (C-term) - Images

TRIM56 Antibody (C-term) (Cat. #AP11813b) western blot analysis in A375 cell line lysates (35ug/lane). This demonstrates the TRIM56 antibody detected the TRIM56 protein (arrow).

TRIM56 Antibody (C-term) - Background

TRIM56 belongs to the TRIM/RBCC family. It contains 2 B box-type zinc fingers and a RING-type zinc finger. There are 3 named isoforms produced by alternative splicing.

TRIM56 Antibody (C-term) - References

Olsen, J.V., et al. Cell 127(3):635-648(2006)

Olsen, J.V., et al. Cell 127(3):635-648(2006)