

CFD Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12089a**Specification**

CFD Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P00746
Other Accession	P51779 , Q3T0A3 , NP_001919.2
Reactivity	Human
Predicted	Bovine, Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	68-99

CFD Antibody (N-term) - Additional Information**Gene ID** 1675**Other Names**

Complement factor D, Adipsin, C3 convertase activator, Properdin factor D, CFD, DF, PFD

Target/Specificity

This CFD antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 68-99 amino acids from the N-terminal region of human CFD.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CFD Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CFD Antibody (N-term) - Protein Information**Name** CFD**Synonyms** DF, PFD

Function Factor D cleaves factor B when the latter is complexed with factor C3b, activating the C3bbb complex, which then becomes the C3 convertase of the alternate pathway. Its function is homologous to that of C1s in the classical pathway.


Cellular Location
Secreted.

CFD Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CFD Antibody (N-term) - Images




CFD Antibody (N-term) (Cat. #AP12089a) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the CFD antibody detected the CFD protein (arrow).




Anti-CFD Antibody (N-term) at 1:2000 dilution + human plasma lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-CFD Antibody (N-term) at 1:2000 dilution + human plasma lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-CFD Antibody (N-term) at 1:2000 dilution + human plasma lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-CFD Antibody (N-term) at 1:2000 dilution + human plasma lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-CFD Antibody (N-term) at 1:2000 dilution + human plasma lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-CFD Antibody (N-term) at 1:2000 dilution + human plasma lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of lysate from human plasma tissue lysate, using CFD Antibody (N-term)(Cat. #AP12089a). AP12089a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



CFD Antibody (N-term) (Cat. #AP12089a)immunohistochemistry analysis in formalin fixed and paraffin embedded human bladder carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of CFD Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

CFD Antibody (N-term) - Background

The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative complement pathway best known for its role in humoral suppression of infectious agents. This protein is also a serine protease that is secreted by adipocytes into the bloodstream. Finally, the encoded protein has a high level of expression in fat, suggesting a role for adipose tissue in immune system biology.

CFD Antibody (N-term) - References

- Bailey, S.D., et al. Diabetes Care (2010) In press :
- Hietaharju, A., et al. Eur. J. Neurol. 17(2):332-334(2010)
- Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)
- Ciprandi, G., et al. Int. Immunopharmacol. 9(12):1460-1463(2009)

Cerhan, J.R., et al. Br. J. Haematol. 145(5):614-623(2009)