

NEU2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12189A

Specification

NEU2 Antibody (N-term) - Product Information

Application	IF, WB, IHC-P-Leica, FC,E
Primary Accession	Q9Y3R4
Other Accession	NP_005374.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	23-50

NEU2 Antibody (N-term) - Additional Information

Gene ID 4759

Other Names

Sialidase-2, Cytosolic sialidase, N-acetyl-alpha-neuraminidase 2, NEU2

Target/Specificity

This NEU2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 23-50 amino acids from the N-terminal region of human NEU2.

Dilution

IF~~1:10~50
WB~~1:1000
IHC-P-Leica~~1:250
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NEU2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NEU2 Antibody (N-term) - Protein Information

Name NEU2

Function Exo-alpha-sialidase that catalyzes the hydrolytic cleavage of the terminal sialic acid (N-acetylneuraminic acid, Neu5Ac) of a glycan moiety in the catabolism of glycolipids, glycoproteins and oligosaccharides (PubMed:[14613940](#), PubMed:[22228546](#)). Recognizes sialyl linkage positions of the glycan moiety as well as the supramolecular organization of the sialoglycoconjugate. Displays preference for alpha- (2->3)-sialylated GD1a and GT1B gangliosides over alpha-(2->8)- sialylated GD1b, in both monomeric forms and micelles. Hydrolyzes monomeric GM1 ganglioside, but has no activity toward the miscellar form (PubMed:[14613940](#)). Has lower sialidase activity for glycoproteins such as fetuin and TF/transferrin that carry a mixture of alpha-(2->3) and alpha-(2->6)-sialyl linkages. Cleaves milk oligosaccharide alpha-(2->3)-sialyllactose, but is inactive toward alpha-(2->6)-sialyllactose isomer. Has no activity toward colominic acid, a homomer of alpha- (2->8)-linked Neu5Ac residues (PubMed:[14613940](#)).

Cellular Location

Cytoplasm, cytosol.


Tissue Location

Expressed in skeletal muscle, fetal liver and embryonic carcinoma cell line NT2-D1.


NEU2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NEU2 Antibody (N-term) - Images


Confocal immunofluorescent analysis of NEU2 Antibody (N-term)(Cat#AP12189a) with A549 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red). DAPI was used to stain the cell nuclear (blue).




Anti-NEU2 Antibody (N-term) at 1:1000 dilution + human fetal liver lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 42 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-NEU2 Antibody (N-term) at 1:2000 dilution + Human heart tissue lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 42 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




NEU2 Antibody (N-term) (Cat. #AP12189a) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the NEU2 antibody detected the NEU2 protein (arrow).



Immunohistochemical analysis of AP12189a on paraffin-embedded human skeletal muscle tissue was performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9.0). Samples were incubated with primary antibody (1:250) for 15min at room temperature. Leica Bond Polymer Refine Detection was used as the secondary antibody.



Immunohistochemical analysis of AP12189a on paraffin-embedded human kidney tissue was performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9.0). Samples were incubated with primary antibody(1:250) for 15min at room temperature. Leica Bond Polymer Refine Detection was used as the secondary antibody.



NEU2 Antibody (N-term) (Cat. #AP12189a) flow cytometric analysis of A549 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

NEU2 Antibody (N-term) - Background

This gene belongs to a family of glycohydrolytic enzymes which remove sialic acid residues from glycoproteins and glycolipids. Expression studies in COS7 cells confirmed that this gene encodes a functional sialidase. Its cytosolic localization was demonstrated by cell fractionation experiments. [provided by RefSeq].

NEU2 Antibody (N-term) - References

- Stoppani, E., et al. Cell Biol. Int. 33(9):1020-1025(2009)
- Li, C.Y., et al. Cell Res. 17(4):357-362(2007)
- Chavas, L.M., et al. J. Biol. Chem. 280(1):469-475(2005)
- Seyrantepe, V., et al. J. Biol. Chem. 279(35):37021-37029(2004)

Tringali, C., et al. J. Biol. Chem. 279(5):3169-3179(2004)