10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CCNYL3 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12215a

Specification

CCNYL3 Antibody (N-term) - Product Information

Application	**WB, FC,E**
Primary Accession	P0C7X3
Other Accession	XP_930958.4
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**39078**
Antigen Region	**46-74**

CCNYL3 Antibody (N-term) - Additional Information

Other Names
Putative cyclin-Y-like protein 3, CCNYL3

Target/Specificity
This CCNYL3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 46-74 amino acids from the N-terminal region of human CCNYL3.

Dilution
WB~~1:1000
FC~~1:10~50

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
CCNYL3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CCNYL3 Antibody (N-term) - Protein Information

Name CCNYL3


CCNYL3 Antibody (N-term) - Protocols

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CCNYL3 Antibody (N-term) - Images



CCNYL3 Antibody (N-term) (Cat. #AP12215a) western blot analysis in NCI-H460 cell line lysates (35ug/lane).This demonstrates the CCNYL3 antibody detected the CCNYL3 protein (arrow).



CCNYL3 Antibody (N-term) (Cat. #AP12215a) flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.