10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

UNQ3029 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12221c

Specification

UNQ3029 Antibody (Center) - Product Information

Application	**WB, FC,E**
Primary Accession	Q6UWF6
Other Accession	XP_001722754.1
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Antigen Region	**62-91**

UNQ3029 Antibody (Center) - Additional Information

Other Names
Putative uncharacterized protein UNQ3029/PRO9830, hypothetical protein

Target/Specificity
This UNQ3029 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 62-91 amino acids from the Central region of human UNQ3029.

Dilution
WB~~1:1000
FC~~1:10~50

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
UNQ3029 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

UNQ3029 Antibody (Center) - Protein Information


UNQ3029 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

UNQ3029 Antibody (Center) - Images



UNQ3029 Antibody (Center) (Cat. #AP12221c) western blot analysis in NCI-H460 cell line lysates (35ug/lane).This demonstrates the UNQ3029 antibody detected the UNQ3029 protein (arrow).



UNQ3029 Antibody (Center) (Cat. #AP12221c) flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.