

OR51Q1 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12319a

Specification

OR51Q1 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q8NH59
Other Accession	NP_001004757.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35747
Antigen Region	34-62

OR51Q1 Antibody (N-term) - Additional Information

Gene ID 390061

Other Names

Olfactory receptor 51Q1, OR51Q1

Target/Specificity

This OR51Q1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 34-62 amino acids from the N-terminal region of human OR51Q1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR51Q1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR51Q1 Antibody (N-term) - Protein Information

Name OR51Q1

Function Odorant receptor.


Cellular Location

Cell membrane; Multi-pass membrane protein.

OR51Q1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR51Q1 Antibody (N-term) - Images

Anti-OR51Q1 Antibody (N-term) at 1:1000 dilution + MDA-MB-453 whole cell lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 36 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

OR51Q1 Antibody (N-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR51Q1 Antibody (N-term) - References

Taylor, T.D., et al. Nature 440(7083):497-500(2006)