

IGSF8 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12389c**Specification**

IGSF8 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	O969P0
Other Accession	NP_443100.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	65034
Antigen Region	226-254

IGSF8 Antibody (Center) - Additional Information**Gene ID** 93185**Other Names**

Immunoglobulin superfamily member 8, IgSF8, CD81 partner 3, Glu-Trp-Ile EWI motif-containing protein 2, EWI-2, Keratinocytes-associated transmembrane protein 4, KCT-4, LIR-D1, Prostaglandin regulatory-like protein, PGRL, CD316, IGSF8, CD81P3, EWI2, KCT4

Target/Specificity

This IGSF8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 226-254 amino acids from the Central region of human IGSF8.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IGSF8 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

IGSF8 Antibody (Center) - Protein Information**Name** IGSF8

Synonyms CD81P3, EWI2, KCT4

Function May play a key role in diverse functions ascribed to CD81 and CD9 such as oocytes fertilization or hepatitis C virus function. May regulate proliferation and differentiation of keratinocytes. May be a negative regulator of cell motility: suppresses T-cell mobility coordinately with CD81, associates with CD82 to suppress prostate cancer cell migration, regulates epidermoid cell reaggregation and motility on laminin-5 with CD9 and CD81 as key linkers. May also play a role on integrin-dependent morphology and motility functions. May participate in the regulation of neurite outgrowth and maintenance of the neural network in the adult brain.

Cellular Location

Cell membrane; Single-pass membrane protein

Tissue Location


Expressed in brain, kidney, testis, liver and placenta with moderate expression in all other tissues. Detected on a majority of B-cells, T-cells, and natural killer cells but not on monocytes, polynuclear cells and platelets

IGSF8 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


IGSF8 Antibody (Center) - Images



Anti-IGSF8 Antibody (Center) at 1:1000 dilution + human brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 65 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



IGSF8 Antibody (Center) (Cat. #AP12389c) western blot analysis in mouse cerebellum tissue lysates (35ug/lane). This demonstrates the IGSF8 antibody detected the IGSF8 protein (arrow).



IGSF8 Antibody (Center) (Cat. #AP12389c) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of IGSF8 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

IGSF8 Antibody (Center) - Background

IGSF8 may play a key role in diverse functions ascribed to CD81 and CD9 such as oocytes fertilization or hepatitis C virus function. May regulate proliferation and differentiation of keratinocytes. May be a negative regulator of cell motility: suppresses T-cell mobility coordinately with CD81, associates with CD82 to suppress prostate cancer cell migration, regulates epidermoid cell reaggregation and motility on laminin-5 with CD9 and CD81 as key linkers. May also play a role on integrin-dependent morphology and motility functions. May participate in the regulation of neurite outgrowth and maintenance of the neural network in the adult brain.

IGSF8 Antibody (Center) - References

Kolesnikova, T.V., et al. Neoplasia 11(1):77-86(2009)
Rocha-Perugini, V., et al. PLoS ONE 3 (4), E1866 (2008) :
Kettner, S., et al. Mol. Cell. Biol. 27(21):7718-7726(2007)

Sala-Valdes, M., et al. J. Biol. Chem. 281(28):19665-19675(2006)

Yang, X.H., et al. J. Biol. Chem. 281(18):12976-12985(2006)