10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

OR8D4 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12537b

Specification

OR8D4 Antibody (C-term) - Product Information

Application	**WB,E**
Primary Accession	Q8NGM9
Other Accession	NP_001005197.1
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**35019**
Antigen Region	**280-308**

OR8D4 Antibody (C-term) - Additional Information

Gene ID 338662

Other Names
Olfactory receptor 8D4, Olfactory receptor OR11-275, OR8D4

Target/Specificity
This OR8D4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 280-308 amino acids from the C-terminal region of human OR8D4.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
OR8D4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR8D4 Antibody (C-term) - Protein Information

Name OR8D4

Function Odorant receptor.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999


Cellular Location
Cell membrane; Multi-pass membrane protein.

OR8D4 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

OR8D4 Antibody (C-term) - Images



OR8D4 Antibody (C-term) (Cat. #AP12537b) western blot analysis in human placenta tissue lysates (35ug/lane).This demonstrates the OR8D4 antibody detected the OR8D4 protein (arrow).

OR8D4 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR8D4 Antibody (C-term) - References

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)