

**PEX14 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP12560c**

**Specification**

---

**PEX14 Antibody (Center) - Product Information**

Application	WB, FC,E
Primary Accession	<a href="#">O75381</a>
Other Accession	<a href="#">NP_004556.1</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	41237
Antigen Region	121-150

**PEX14 Antibody (Center) - Additional Information**

**Gene ID** 5195

**Other Names**

Peroxisomal membrane protein PEX14, PTS1 receptor-docking protein, Peroxin-14, Peroxisomal membrane anchor protein PEX14, PEX14

**Target/Specificity**

This PEX14 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 121-150 amino acids from the Central region of human PEX14.

**Dilution**

WB~~1:1000  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PEX14 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**PEX14 Antibody (Center) - Protein Information**

**Name** PEX14 {ECO:0000303|PubMed:9653144, ECO:0000312|HGNC:HGNC:8856}

**Function** Component of the PEX13-PEX14 docking complex, a translocon channel that specifically mediates the import of peroxisomal cargo proteins bound to PEX5 receptor (PubMed:[9653144](#), PubMed:[24235149](#), PubMed:[28765278](#)). The PEX13-PEX14 docking complex forms a large import pore which can be opened to a diameter of about 9 nm (By similarity). Mechanistically, PEX5 receptor along with cargo proteins associates with the PEX14 subunit of the PEX13-PEX14 docking complex in the cytosol, leading to the insertion of the receptor into the organelle membrane with the concomitant translocation of the cargo into the peroxisome matrix (PubMed:[24235149](#), PubMed:[28765278](#)). Plays a key role for peroxisome movement through a direct interaction with tubulin (PubMed:[21525035](#)).

#### Cellular Location


Peroxisome membrane; Single-pass membrane protein {ECO:0000250|UniProtKB:Q642G4}

#### PEX14 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### PEX14 Antibody (Center) - Images



PEX14 Antibody (Center) (Cat. #AP12560c) western blot analysis in mouse Neuro-2a cell line lysates (35ug/lane). This demonstrates the PEX14 antibody detected the PEX14 protein (arrow).



PEX14 Antibody (Center) (Cat. #AP12560c) flow cytometric analysis of Neuro-2a cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated donkey-anti-rabbit secondary antibodies were used for the analysis.

#### **PEX14 Antibody (Center) - Background**

This gene encodes an essential component of the peroxisomal import machinery. The protein is integrated into peroxisome membranes with its C-terminus exposed to the cytosol, and interacts with the cytosolic receptor for proteins containing a PTS1 peroxisomal targeting signal. The protein also functions as a transcriptional corepressor and interacts with a histone deacetylase. A mutation in this gene results in one form of Zellweger syndrome.

#### **PEX14 Antibody (Center) - References**

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010) Macgregor, S., et al. Hum. Mol. Genet. 19(13):2716-2724(2010) Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) : Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009) Shiozawa, K., et al. J. Biol. Chem. 284(37):25334-25342(2009)