10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

EBNA1BP2 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12662b

Specification

EBNA1BP2 Antibody (C-term) - Product Information

Application	**WB,E**
Primary Accession	Q99848
Other Accession	NP_001153408.1, NP_006815.2
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**34852**
Antigen Region	**271-302**

EBNA1BP2 Antibody (C-term) - Additional Information

Gene ID 10969

Other Names
Probable rRNA-processing protein EBP2, EBNA1-binding protein 2, Nucleolar protein p40, EBNA1BP2, EBP2

Target/Specificity
This EBNA1BP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 271-302 amino acids from the C-terminal region of human EBNA1BP2.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
EBNA1BP2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

EBNA1BP2 Antibody (C-term) - Protein Information

Name EBNA1BP2

Synonyms EBP2

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Function Required for the processing of the 27S pre-rRNA.

Cellular Location
Nucleus, nucleolus. Note=Associated with the nucleolus in an RNA-dependent manner


Tissue Location
Ubiquitous..

EBNA1BP2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

EBNA1BP2 Antibody (C-term) - Images



EBNA1BP2 Antibody (C-term) (Cat. #AP12662b) western blot analysis in NCI-H292 cell line lysates (35ug/lane).This demonstrates the EBNA1BP2 antibody detected the EBNA1BP2 protein (arrow).

EBNA1BP2 Antibody (C-term) - Background

EBNA1BP2 is required for the processing of the 27S pre-rRNA (By similarity).