

PCDHB11 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12988a

Specification

PCDHB11 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	<u>Q9Y5F2</u>
Other Accession	<u>NP_061754.1</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	87088
Antigen Region	47-76

PCDHB11 Antibody (N-term) - Additional Information

Gene ID 56125

Other Names Protocadherin beta-11, PCDH-beta-11, PCDHB11

Target/Specificity This PCDHB11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 47-76 amino acids from the N-terminal region of human PCDHB11.

Dilution WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions PCDHB11 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PCDHB11 Antibody (N-term) - Protein Information

Name PCDHB11

Function Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.



Cellular Location


Cell membrane; Single-pass type I membrane protein

PCDHB11 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

PCDHB11 Antibody (N-term) - Images



PCDHB11 Antibody (N-term) (Cat. #AP12988a) western blot analysis in NCI-H292 cell line lysates (35ug/lane).This demonstrates the PCDHB11 antibody detected the PCDHB11 protein (arrow).

PCDHB11 Antibody (N-term) - Background

This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the transcripts from these genes are made up of only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific cell-cell neural connections.

PCDHB11 Antibody (N-term) - References



Gregorio, S.P., et al. Psychiatry Res 165 (1-2), 1-9 (2009) : Frank, M., et al. Curr. Opin. Cell Biol. 14(5):557-562(2002) Vanhalst, K., et al. FEBS Lett. 495 (1-2), 120-125 (2001) : Wu, Q., et al. Genome Res. 11(3):389-404(2001) Nollet, F., et al. J. Mol. Biol. 299(3):551-572(2000)