10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Bik BH3 Domain Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1319a

Specification

Bik BH3 Domain Antibody - Product Information

Application	**WB, IHC-P,E**
Primary Accession	Q13323
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Antigen Region	**39-74**

Bik BH3 Domain Antibody - Additional Information

Gene ID 638

Other Names
Bcl-2-interacting killer, Apoptosis inducer NBK, BIP1, BP4, BIK, NBK

Target/Specificity
This Bik BH3 Domain antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 39-74 amino acids from human Bik BH3 Domain.

Dilution
WB~~1:2000
IHC-P~~1:50~100

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
Bik BH3 Domain Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Bik BH3 Domain Antibody - Protein Information

Name BIK

Synonyms NBK

Function Accelerates programmed cell death. Association to the apoptosis repressors Bcl-X(L),

BHRF1, Bcl-2 or its adenovirus homolog E1B 19k protein suppresses this death-promoting activity. Does not interact with BAX.


Cellular Location
Endomembrane system; Single-pass membrane protein. Mitochondrion membrane; Single-pass membrane protein. Note=Around the nuclear envelope, and in cytoplasmic membranes

Bik BH3 Domain Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Bik BH3 Domain Antibody - Images




Bik Antibody (BH3) (Cat. #AP1319a) western blot analysis in A549 cell line lysates (35ug/lane).This demonstrates the Bik antibody detected the Bik protein (arrow).

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



All lanes : Anti-hBik-BH3 at 1:2000 dilution Lane 1: A431 whole cell lysate Lane 2: Ramos whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 18 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Bik BH3 Domain Antibody - Background

The Bik protein is known to interact with cellular and viral survival-promoting proteins, such as BCL2 and the Epstein-Barr virus in order to enhance programmed cell death. Because its activity is suppressed in the presence of survival-promoting proteins, this protein is suggested as a likely target for antiapoptotic proteins. This protein shares a critical BH3 domain with other death-promoting proteins, BAX and BAK.

Bik BH3 Domain Antibody - References

Arena, V., et al., Genes Chromosomes Cancer 38(1):91-96 (2003).
Gillissen, B., et al., EMBO J. 22(14):3580-3590 (2003).
Germain, M., et al., J. Biol. Chem. 277(20):18053-18060 (2002).
Zou, Y., et al., Cancer Res. 62(1):8-12 (2002).
Castells, A., et al., Gastroenterology 117(4):831-837 (1999).

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Bik BH3 Domain Antibody - Citations

- TNF pretreatment interferes with mitochondrial apoptosis in the mouse liver by A20-mediated down-regulation of Bax.