

CASP10 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1326c**Specification**

CASP10 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O92851
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	58951
Antigen Region	164-192

CASP10 Antibody (Center) - Additional Information**Gene ID** 843**Other Names**

Caspase-10, CASP-10, Apoptotic protease Mch-4, FAS-associated death domain protein interleukin-1B-converting enzyme 2, FLICE2, ICE-like apoptotic protease 4, Caspase-10 subunit p23/17, Caspase-10 subunit p12, CASP10, MCH4

Target/Specificity

This CASP10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 164-192 amino acids from the Central region of human CASP10.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CASP10 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CASP10 Antibody (Center) - Protein Information**Name** CASP10**Synonyms** MCH4

Function Involved in the activation cascade of caspases responsible for apoptosis execution. Recruited to both Fas- and TNFR-1 receptors in a FADD dependent manner. May participate in the granzyme B apoptotic pathways. Cleaves and activates effector caspases CASP3, CASP4, CASP6, CASP7, CASP8 and CASP9. Hydrolyzes the small- molecule substrates, Tyr- Val-Ala-Asp-|-AMC and Asp-Glu-Val-Asp-|-AMC.

Tissue Location


Detectable in most tissues. Lowest expression is seen in brain, kidney, prostate, testis and colon

CASP10 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CASP10 Antibody (Center) - Images



Western blot analysis of CASP10 antibody (Center) (Cat.#AP1326c) in HepG2 cell line lysates (35ug/lane). CASP10 (arrow) was detected using the purified Pab.

CASP10 Antibody (Center) - Background

CASP10 is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 3 and 7, and the protein itself is processed by caspase 8. Mutations in the protein are associated with apoptosis defects seen in type II autoimmune lymphoproliferative syndrome.

CASP10 Antibody (Center) - References

Lan,Q., Morton,L.M. Blood 114 (2), 264-267 (2009)

Kim,Y.R., Kim,K.M. Hum. Pathol. 40 (6), 868-871 (2009)

Lisa-Santamaria,P. Biochim. Biophys. Acta 1793 (3), 561-571 (2009)