

CHMP2B Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13303a**Specification**

CHMP2B Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O9UQN3
Other Accession	O8BJF9 , O3SX42 , NP_054762.2
Reactivity	Human
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	23907
Antigen Region	1-30

CHMP2B Antibody (N-term) - Additional Information**Gene ID** 25978**Other Names**

Charged multivesicular body protein 2b, CHMP25, Chromatin-modifying protein 2b, CHMP2b, Vacuolar protein sorting-associated protein 2-2, Vps2-2, hVps2-2, CHMP2B

Target/Specificity

This CHMP2B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human CHMP2B.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CHMP2B Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CHMP2B Antibody (N-term) - Protein Information**Name** CHMP2B

Function Probable core component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I, -II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis and the budding of enveloped viruses (HIV-1 and other lentiviruses). ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4.

Cellular Location

Cytoplasm, cytosol. Late endosome membrane; Peripheral membrane protein

Tissue Location


Widely expressed. Expressed in brain, heart, skeletal muscle, spleen, kidney, liver, small intestine, pancreas, lung, placenta and leukocytes. In brain, it is expressed in cerebellum, cerebral cortex, medulla, spinal cord, occipital lobe, frontal lobe, temporal lobe and putamen.

CHMP2B Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CHMP2B Antibody (N-term) - Images



CHMP2B Antibody (N-term) (Cat. #AP13303a) western blot analysis in SK-BR-3 cell line lysates

(35ug/lane). This demonstrates the CHMP2B antibody detected the CHMP2B protein (arrow).



CHMP2B Antibody (N-term) (Cat. #AP13303a) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CHMP2B Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

CHMP2B Antibody (N-term) - Background

This gene encodes a component of the heteromeric ESCRT-III complex (Endosomal Sorting Complex Required for Transport III) that functions in the recycling or degradation of cell surface receptors. ESCRT-III functions in the concentration and invagination of ubiquitinated endosomal cargos into intraluminal vesicles. The protein encoded by this gene is found as a monomer in the cytosol or as an oligomer in ESCRT-III complexes on endosomal membranes. It is expressed in neurons of all major regions of the brain. Mutations in this gene result in one form of familial frontotemporal lobar degeneration.

CHMP2B Antibody (N-term) - References

Kaivorinne, A.L., et al. Eur. J. Neurol. 17(11):1393-1395(2010)
Ghanim, M., et al. J. Neurol. (2010) In press :
Yamazaki, Y., et al. Neurosci. Lett. 477(2):86-90(2010)
Tsai, C.P., et al. Neurobiol. Aging (2010) In press :
Cox, L.E., et al. PLoS ONE 5 (3), E9872 (2010) :