

CACNA2D2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13380C**Specification**

CACNA2D2 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q9NY47
Other Accession	NP_001167522.1 , NP_001005505.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	643-671

CACNA2D2 Antibody (Center) - Additional Information**Gene ID** 9254**Other Names**

Voltage-dependent calcium channel subunit alpha-2/delta-2, Voltage-gated calcium channel subunit alpha-2/delta-2, Voltage-dependent calcium channel subunit alpha-2-2, Voltage-dependent calcium channel subunit delta-2, CACNA2D2, KIAA0558

Target/Specificity

This CACNA2D2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 643-671 amino acids from the Central region of human CACNA2D2.

Dilution

WB~~1:2000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CACNA2D2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CACNA2D2 Antibody (Center) - Protein Information**Name** CACNA2D2

Synonyms KIAA0558

Function The alpha-2/delta subunit of voltage-dependent calcium channels regulates calcium current density and activation/inactivation kinetics of the calcium channel. Acts as a regulatory subunit for P/Q- type calcium channel (CACNA1A), N-type (CACNA1B), L-type (CACNA1C OR CACNA1D) and possibly T-type (CACNA1G) (PubMed:[15111129](#), PubMed:[23339110](#)). Overexpression induces apoptosis.

Cellular Location

Membrane; Single-pass type I membrane protein. Note=Colocalizes with CACNA1A in lipid raft fractions.


Tissue Location

Predominantly present in cerebellar cortex. Present in various lung tumor cell lines, while it is absent in normal lung (at protein level). Highly expressed in heart, lung, testis, pancreas and skeletal muscle. Also expressed in kidney, liver, placenta and brain


CACNA2D2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CACNA2D2 Antibody (Center) - Images

Anti-CACNA2D2 Antibody (Center) at 1:2000 dilution + human testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 130 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-CACNA2D2 Antibody (Center) at 1:2000 dilution + mouse kidney lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 130 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



CACNA2D2 Antibody (Center) (Cat. #AP13380c) immunohistochemistry analysis in formalin fixed and paraffin embedded human cerebellum tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CACNA2D2 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

CACNA2D2 Antibody (Center) - Background

This gene encodes a member of the alpha-2/delta subunit family, a protein in the voltage-dependent calcium channel complex. Calcium channels mediate the influx of calcium ions into the cell upon membrane polarization and consist of a complex of alpha-1, alpha-2/delta, beta, and gamma subunits in a 1:1:1:1 ratio. Various versions of each of these subunits exist, either expressed from similar genes or the result of alternative splicing. Research on a highly similar protein in rabbit suggests the protein described in this record is cleaved into alpha-2 and delta subunits. Alternate transcriptional splice variants of this gene, encoding different isoforms, have been characterized.

CACNA2D2 Antibody (Center) - References

Donato, R., et al. J. Neurosci. 26(48):12576-12586(2006) Ivanov, S.V., et al. Am. J. Pathol. 165(3):1007-1018(2004) Carboni, G.L., et al. Oncogene 22(4):615-626(2003) Honorio, S., et al. Mol. Cell. Probes 15(6):391-393(2001) Angeloni, D., et al. Mol. Cell. Probes 15(2):125-127(2001)

CACNA2D2 Antibody (Center) - Citations

- [Discovery of Biomarker Panels for Neural Dysfunction in Inborn Errors of Amino Acid Metabolism.](#)