

PCDH18 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP13587B

Specification

PCDH18 Antibody (C-term) - Product Information

Application WB,E **Primary Accession** O9HCL0 NP 061908.1 Other Accession Reactivity Mouse Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 126149 Antigen Region 970-999

PCDH18 Antibody (C-term) - Additional Information

Gene ID 54510

Other Names

Protocadherin-18, PCDH18, KIAA1562

Target/Specificity

This PCDH18 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 970-999 amino acids from the C-terminal region of human PCDH18.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PCDH18 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PCDH18 Antibody (C-term) - Protein Information

Name PCDH18

Synonyms KIAA1562



Function Potential calcium-dependent cell-adhesion protein.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location


Expressed in all tissues, with highest expression in lung and ovary.

PCDH18 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture


PCDH18 Antibody (C-term) - Images



PCDH18 Antibody (C-term) (Cat. #AP13587b) western blot analysis in mouse liver tissue lysates (35ug/lane). This demonstrates the PCDH18 antibody detected the PCDH18 protein (arrow).

PCDH18 Antibody (C-term) - Background

This gene belongs to the protocadherin gene family, a subfamily of the cadherin superfamily. This gene encodes a protein which contains 6 extracellular cadherin domains, a transmembrane domain and a cytoplasmic tail differing from those of the classical cadherins. Although its specific function is undetermined, the cadherin-related neuronal receptor is thought to play a role in the establishment and function of specific cell-cell connections in the brain.





PCDH18 Antibody (C-term) - References

Homayouni, R., et al. Biochem. Biophys. Res. Commun. 289(2):539-547(2001) Wolverton, T., et al. Genomics 76 (1-3), 66-72 (2001): Suzuki, S.T. Exp. Cell Res. 261(1):13-18(2000) Nollet, F., et al. J. Mol. Biol. 299(3):551-572(2000) Yagi, T., et al. Genes Dev. 14(10):1169-1180(2000)