

ADAL Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP13621b

Specification

ADAL Antibody (C-term) - Product Information

Application WB, IHC-P,E
Primary Accession Q6DHV7

Other Accession NP 001012987.1, NP 001152752.1

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Mouse
Rabbit
Polyclonal
Rabbit IgG
40264
317-344

ADAL Antibody (C-term) - Additional Information

Gene ID 161823

Other Names

Adenosine deaminase-like protein, 354-, ADAL

Target/Specificity

This ADAL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 317-344 amino acids from the C-terminal region of human ADAL.

Dilution

WB~~1:1000 IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ADAL Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ADAL Antibody (C-term) - Protein Information

Name ADAL

Synonyms ADAL1 {ECO:0000303|PubMed:21755941}




Function Catalyzes the hydrolysis of the free cytosolic methylated adenosine nucleotide N(6)-methyl-AMP (N6-mAMP) to produce inositol monophosphate (IMP) and methylamine (PubMed:21755941, PubMed:29884623). Is required for the catabolism of cytosolic N6-mAMP, which is derived from the degradation of mRNA containing N6-methylated adenine (m6A) (PubMed:21755941, PubMed:29884623). Catalyzes the removal of different alkyl groups not only from N6-substituted purine or 2-aminopurine nucleoside monophosphates but also from O6-substituted compounds in vitro (PubMed:21755941).

ADAL Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

ADAL Antibody (C-term) - Images



ADAL Antibody (C-term) (Cat. #AP13621b) western blot analysis in mouse kidney,heart tissue lysates (35ug/lane). This demonstrates the ADAL antibody detected the ADAL protein (arrow).





ADAL Antibody (C-term) (Cat. #AP13621b)immunohistochemistry analysis in formalin fixed and paraffin embedded human esophagus carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of ADAL Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

ADAL Antibody (C-term) - Background

ADAL is a putative nucleoside deaminase. It may catalyze the hydrolytic deamination of adenosine or some similar substrate and play a role in purine metabolism (By similarity).

ADAL Antibody (C-term) - References

Strausberg, R.L., et al. Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903(2002)