10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SLC2A13 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13677C

Specification

SLC2A13 Antibody (Center) - Product Information

Application	**WB,E**
Primary Accession	Q96QE2
Other Accession	NP_443117.3
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**70371**
Antigen Region	**268-297**

SLC2A13 Antibody (Center) - Additional Information

Gene ID 114134

Other Names
Proton myo-inositol cotransporter, H(+)-myo-inositol cotransporter, Hmit, H(+)-myo-inositol symporter, Solute carrier family 2 member 13, SLC2A13

Target/Specificity
This SLC2A13 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 268-297 amino acids from the Central region of human SLC2A13.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
SLC2A13 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

SLC2A13 Antibody (Center) - Protein Information

Name SLC2A13 (HGNC:15956)

Function H(+)-myo-inositol cotransporter (PubMed:11500374). Can also transport related

stereoisomers (PubMed:11500374).

Cellular Location
Cell membrane; Multi-pass membrane protein


Tissue Location
Predominantly expressed in the brain.

SLC2A13 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

SLC2A13 Antibody (Center) - Images



Anti-SLC2A13 Antibody (Center) at 1:1000 dilution + human brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 70 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

SLC2A13 Antibody (Center) - Background

H(+)-myo-inositol cotransporter.It can also transport related stereoisomers.

SLC2A13 Antibody (Center) - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) : Bankovic, J., et al. Lung Cancer 67(2):151-159(2010) Satake, W., et al. Nat. Genet. 41(12):1303-1307(2009) Di Daniel, E., et al. BMC Cell Biol. 10, 54 (2009) : Barrett, J.C., et al. Nat. Genet. 40(8):955-962(2008)