

PPP2CB Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13765b**Specification**

PPP2CB Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P62714
Other Accession	P23696 , P62716 , P11611 , P11493 , P62715 , Q0P594 , P63331 , P67777 , P67776 , P63330 , P67775 , P48463 , P67774 , NP_001009552.1 , G5EGK8
Reactivity	Human
Predicted	Bovine, Chicken, Mouse, Pig, Rabbit, Rat, C.Elegans, Drosophila
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35575
Antigen Region	278-307

PPP2CB Antibody (C-term) - Additional Information**Gene ID** 5516**Other Names**

Serine/threonine-protein phosphatase 2A catalytic subunit beta isoform, PP2A-beta, PPP2CB

Target/Specificity

This PPP2CB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 278-307 amino acids from the C-terminal region of human PPP2CB.

Dilution

WB~~1:1000

IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PPP2CB Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PPP2CB Antibody (C-term) - Protein Information

Name PPP2CB ([HGNC:9300](#))

Function Catalytic subunit of protein phosphatase 2A (PP2A), a serine/threonine phosphatase involved in the regulation of a wide variety of enzymes, signal transduction pathways, and cellular events (Probable). PP2A can modulate the activity of phosphorylase B kinase, casein kinase 2, mitogen-stimulated S6 kinase, and MAP-2 kinase.

Cellular Location


Cytoplasm. Nucleus. Chromosome, centromere. Cytoplasm, cytoskeleton, spindle pole. Note=In prometaphase cells, but not in anaphase cells, localizes at centromeres. During mitosis, also found at spindle poles

PPP2CB Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PPP2CB Antibody (C-term) - Images



PPP2CB Antibody (C-term) (Cat. #AP13765b) western blot analysis in T47D cell line lysates (35ug/lane). This demonstrates the PPP2CB antibody detected the PPP2CB protein (arrow).



PPP2CB Antibody (C-term) (AP13765b) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of PPP2CB Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

PPP2CB Antibody (C-term) - Background

PP2A can modulate the activity of phosphorylase B kinase casein kinase 2, mitogen-stimulated S6 kinase, and MAP-2 kinase.