

**OR1N1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP13812b****Specification**

---

**OR1N1 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q8NGS0</a>
Other Accession	<a href="#">NP_036495.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	34650
Antigen Region	277-306

**OR1N1 Antibody (C-term) - Additional Information****Gene ID** 138883**Other Names**

Olfactory receptor 1N1, Olfactory receptor 1-26, OR1-26, Olfactory receptor 1N3, Olfactory receptor OR9-22, OR1N1, OR1N3

**Target/Specificity**

This OR1N1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 277-306 amino acids from the C-terminal region of human OR1N1.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

OR1N1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**OR1N1 Antibody (C-term) - Protein Information****Name** OR1N1**Synonyms** OR1N3

**Function** Odorant receptor.

**Cellular Location**


Cell membrane; Multi-pass membrane protein.

**OR1N1 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**OR1N1 Antibody (C-term) - Images**



OR1N1 Antibody (C-term) (Cat. #AP13812b) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the OR1N1 antibody detected the OR1N1 protein (arrow).

**OR1N1 Antibody (C-term) - Background**

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

**OR1N1 Antibody (C-term) - References**

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)

Fuchs, T., et al. Genomics 80(3):295-302(2002)  
Rouquier, S., et al. Nat. Genet. 18(3):243-250(1998)