

PIAS4 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14013a

Specification

PIAS4 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q8N2W9
Other Accession	NP_056981.2
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	56504
Antigen Region	44-73

PIAS4 Antibody (N-term) - Additional Information

Gene ID 51588

Other Names

E3 SUMO-protein ligase PIAS4, 632-, PIASy, Protein inhibitor of activated STAT protein 4, Protein inhibitor of activated STAT protein gamma, PIAS-gamma, PIAS4, PIASG

Target/Specificity

This PIAS4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 44-73 amino acids from the N-terminal region of human PIAS4.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PIAS4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PIAS4 Antibody (N-term) - Protein Information

Name PIAS4 {ECO:0000303|PubMed:32832608, ECO:0000312|HGNC:HGNC:17002}

Function Functions as an E3-type small ubiquitin-like modifier (SUMO) ligase, stabilizing the

interaction between UBE2I and the substrate, and as a SUMO-tethering factor (PubMed:[12511558](#), PubMed:[12631292](#), PubMed:[12727872](#), PubMed:[15831457](#), PubMed:[15976810](#), PubMed:[22508508](#), PubMed:[32832608](#)). Mediates sumoylation of CEBPA, PARK7, HERC2, MYB, TCF4 and RNF168 (PubMed:[12511558](#), PubMed:[12631292](#), PubMed:[12727872](#), PubMed:[15831457](#), PubMed:[15976810](#), PubMed:[22508508](#)). Plays a crucial role as a transcriptional coregulation in various cellular pathways, including the STAT pathway, the p53/TP53 pathway, the Wnt pathway and the steroid hormone signaling pathway (PubMed:[11388671](#)). Involved in gene silencing (PubMed:[11248056](#)). In Wnt signaling, represses LEF1 and enhances TCF4 transcriptional activities through promoting their sumoylations (PubMed:[12727872](#), PubMed:[15831457](#)). Enhances the sumoylation of MTA1 and may participate in its paralog-selective sumoylation (PubMed:[21965678](#)). Binds to AT-rich DNA sequences, known as matrix or scaffold attachment regions (MARs/SARs) (By similarity). Catalyzes conjugation of SUMO2 to KAT5 in response to DNA damage, facilitating repair of DNA double-strand breaks (DSBs) via homologous recombination (HR) (PubMed:[32832608](#)). Mediates sumoylation of PARP1 in response to PARP1 trapping to chromatin (PubMed:[35013556](#)).

Cellular Location

Nucleus, PML body Note=Colocalizes with SUMO1 and TCF7L2/TCF4 and LEF1 in a subset of PML (promyelocytic leukemia) nuclear bodies.

Tissue Location


Highly expressed in testis and, at lower levels, in spleen, prostate, ovary, colon and peripheral blood leukocytes

PIAS4 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PIAS4 Antibody (N-term) - Images



PIAS4 Antibody (N-term) (Cat. #AP14013a) western blot analysis in mouse heart tissue lysates

(35ug/lane). This demonstrates the PIAS4 antibody detected the PIAS4 protein (arrow).

PIAS4 Antibody (N-term) - Background

PIAS4 functions as an E3-type small ubiquitin-like modifier (SUMO) ligase, stabilizing the interaction between UBE2I and the substrate, and as a SUMO-tethering factor. Plays a crucial role as a transcriptional coregulation in various cellular pathways, including the STAT pathway, the p53 pathway, the Wnt pathway and the steroid hormone signaling pathway. Involved in gene silencing. Promotes PARK7 sumoylation. In Wnt signaling, represses LEF1 and enhances TCF4 transcriptional activities through promoting their sumoylations.

PIAS4 Antibody (N-term) - References

Kang, X., et al. *Oncogene* 29(41):5568-5578(2010)
Liu, Y., et al. *J. Invest. Dermatol.* 130(5):1384-1390(2010)
Zhang, C., et al. *Cell. Immunol.* 263(2):235-240(2010)
Wang, J., et al. *J. Biol. Chem.* 283(34):23235-23243(2008)
Imoto, S., et al. *Biochem. Biophys. Res. Commun.* 370(2):359-365(2008)