

RDX Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP14028B

Specification

RDX Antibody (C-term) - Product Information

Application WB,E
Primary Accession P35241

Other Accession <u>P26044</u>, <u>Q9PU45</u>, <u>Q32LP2</u>, <u>NP_002897.1</u>

Reactivity Human

Predicted Bovine, Chicken, Pig

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 68564
Antigen Region 466-495

RDX Antibody (C-term) - Additional Information

Gene ID 5962

Other Names

Radixin, RDX

Target/Specificity

This RDX antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 466-495 amino acids from the C-terminal region of human RDX.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RDX Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RDX Antibody (C-term) - Protein Information

Name RDX

Function Probably plays a crucial role in the binding of the barbed end of actin filaments to the



plasma membrane.

Cellular Location


Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytoskeleton. Cleavage furrow. Cell projection, microvillus {ECO:0000250|UniProtKB:P26043}. Note=Highly concentrated in the undercoat of the cell-to-cell adherens junction and the cleavage furrow in the interphase and mitotic phase, respectively

RDX Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

RDX Antibody (C-term) - Images




RDX Antibody (C-term) (Cat. #AP14028b) western blot analysis in U251 cell line lysates (35ug/lane). This demonstrates the RDX antibody detected the RDX protein (arrow).

RDX Antibody (C-term) - Background

Radixin is a cytoskeletal protein that may be important in linking actin to the plasma membrane. It is highly similar in sequence to both ezrin and moesin. The radixin gene has been localized by fluorescence in situ hybridization to 11q23. A truncated version representing a pseudogene (RDXP2) was assigned to Xp21.3. Another pseudogene that seemed to lack introns (RDXP1) was mapped to 11p by Southern and PCR analyses.

RDX Antibody (C-term) - References

Gloerich, M., et al. Mol. Cell. Biol. 30(22):5421-5431(2010) Kramer, B., et al. Eur. J. Immunol. 39(12):3447-3458(2009)





Parisiadou, L., et al. J. Neurosci. 29(44):13971-13980(2009) Belkina, N.V., et al. Proc. Natl. Acad. Sci. U.S.A. 106(12):4707-4712(2009) Wolf, C., et al. BMC Med. Genet. 10, 91 (2009) :