

GRIA3 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14041a**Specification**

GRIA3 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P42263
Other Accession	P19492 , Q9Z2W9 , Q38PU6 , NP_015564.4 , NP_000819.3 , Q71E60
Reactivity	Human
Predicted	Zebrafish, Monkey, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	101157
Antigen Region	188-217

GRIA3 Antibody (N-term) - Additional Information**Gene ID** 2892**Other Names**

Glutamate receptor 3, GluR-3, AMPA-selective glutamate receptor 3, GluR-C, GluR-K3, Glutamate receptor ionotropic, AMPA 3, GluA3, GRIA3, GLUR3, GLURC

Target/Specificity

This GRIA3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 188-217 amino acids from the N-terminal region of human GRIA3.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GRIA3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

GRIA3 Antibody (N-term) - Protein Information**Name** GRIA3

Synonyms GLUR3, GLURC

Function Receptor for glutamate that functions as a ligand-gated ion channel in the central nervous system and plays an important role in excitatory synaptic transmission. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. In the presence of CACNG4 or CACNG7 or CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of glutamate.


Cellular Location

Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane; Multi-pass membrane protein Note=Interaction with CNIH2 and CNIH3 promotes cell surface expression

GRIA3 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GRIA3 Antibody (N-term) - Images

GRIA3 Antibody (N-term) (Cat. #AP14041a) western blot analysis in WiDr cell line lysates (35ug/lane). This demonstrates the GRIA3 antibody detected the GRIA3 protein (arrow).

GRIA3 Antibody (N-term) - Background

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. These receptors are heteromeric protein complexes composed of multiple subunits, arranged to form ligand-gated ion channels. The classification of

glutamate receptors is based on their activation by different pharmacologic agonists. The subunit encoded by this gene belongs to a family of AMPA (alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate)-sensitive glutamate receptors, and is subject to RNA editing (AGA->GGA; R->G). Alternative splicing at this locus results in different isoforms, which may vary in their signal transduction properties.

GRIA3 Antibody (N-term) - References

Ripka, S., et al. Neoplasia 12(8):659-667(2010)
Liu, Q., et al. Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi 26(4):376-378(2010)
Feyissa, A.M., et al. Prog. Neuropsychopharmacol. Biol. Psychiatry 34(2):279-283(2010)
Marek, G.J., et al. Mol. Pharmacol. 77(3):317-326(2010)
Formicola, D., et al. BMC Med. Genet. 11, 103 (2010) :