

DPPA4 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1438a

Specification

DPPA4 Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O7L190
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	33541
Antigen Region	50-79

DPPA4 Antibody (N-term) - Additional Information

Gene ID 55211

Other Names

Developmental pluripotency-associated protein 4, DPPA4

Target/Specificity

This DPPA4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 50-79 amino acids from the N-terminal region of human DPPA4.

Dilution

WB~~1:1000

IHC-P~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DPPA4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DPPA4 Antibody (N-term) - Protein Information

Name DPPA4

Function May be involved in the maintenance of active epigenetic status of target genes. May inhibit differentiation of embryonic cells into a primitive ectoderm lineage.


Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q8CCG4}. Note=Associated with transcriptionally active chromatin {ECO:0000250|UniProtKB:Q8CCG4}


DPPA4 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DPPA4 Antibody (N-term) - Images

Anti-DPPA4 Antibody (N-term) at 1:2000 dilution + NCCIT whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 34 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Western blot analysis of DPPA4 Antibody (N-term) (Cat. #AP1438a) in CEM cell line lysates (35ug/lane). DPPA4 (arrow) was detected using the purified Pab.



AP1438a staining DPPA4 in human testis tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

DPPA4 Antibody (N-term) - Background

DppA4 is a member of the Dppa family of developmental pluripotency associated transcription factors. Dppa4 is a target of Oct4, Sox2 and Nanog. It may play a role in maintaining cell pluripotentiality.

DPPA4 Antibody (N-term) - References

Bortvin, A., Development 130 (8), 1673-1680 (2003)