

POTE Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1439d

Specification

POTE Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q86YR6
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	542-571

POTE Antibody (C-term) - Additional Information

Gene ID 100288966;317754

Other Names

POTE ankyrin domain family member D, ANKRD26-like family B member 3, Ankyrin repeat domain-containing protein 21, Prostate, ovary, testis-expressed protein, Protein POTE, POTED, A26B3, ANKRD21, POTE

Target/Specificity

This POTE antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 542-571 amino acids from the C-terminal region of human POTE.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

POTE Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

POTE Antibody (C-term) - Protein Information

Name POTED

Synonyms A26B3, ANKRD21, POTE

Cellular Location

Cell membrane; Peripheral membrane protein


Tissue Location

Expressed in prostate, ovary, testis, placenta and prostate cancer cell lines. Localizes to basal and terminal prostate epithelial cells.


POTE Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

POTE Antibody (C-term) - Images

Western blot analysis of POTE Antibody (C-term) (Cat. #AP1439d) in Ramos cell line lysates (35ug/lane). POTE (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human prostata carcinoma tissue reacted with POTE

antibody (C-term) (Cat.#AP1439d), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

POTE Antibody (C-term) - Background

The POTE protein contains seven ankyrin repeats between amino acids 140 and 380. Expression of POTE in prostate cancer and its undetectable expression in normal essential tissues make POTE a candidate for the immunotherapy of prostate cancer. The existence of a large number of closely related but rapidly diverging members, their location on multiple chromosomes and their limited expression pattern suggest an important role for the POTE gene family in reproductive processes.

POTE Antibody (C-term) - References

Bera T.K., Proc. Natl. Acad. Sci. U.S.A. 99:16975-16980(2002).
Bera,T.K., Gene 337, 45-53 (2004)