

NUDCD1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP14736c

Specification

NUDCD1 Antibody (Center) - Product Information

Application WB,E
Primary Accession O96RS6

Other Accession NP 116258.2, NP 001121683.1

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Rabbit
Polyclonal
Rabbit IgG
66756
194-223

NUDCD1 Antibody (Center) - Additional Information

Gene ID 84955

Other Names

NudC domain-containing protein 1, Chronic myelogenous leukemia tumor antigen 66, Tumor antigen CML66, NUDCD1 (HGNC:24306)

Target/Specificity

This NUDCD1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 194-223 amino acids from the Central region of human NUDCD1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NUDCD1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

NUDCD1 Antibody (Center) - Protein Information

Name NUDCD1 (HGNC:24306)



Cellular Location

[Isoform 1]: Cytoplasm. Nucleus. [Isoform 3]: Cytoplasm. Nucleus.

Tissue Location


Isoform 1 is specifically expressed in leukemias and a variety of solid tumor cell lines and is also detected in testis and heart. Isoform 2 is predominantly expressed in testis and weakly expressed in tumor cells.

NUDCD1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

NUDCD1 Antibody (Center) - Images



NUDCD1 Antibody (Center) (Cat. #AP14736c) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the NUDCD1 antibody detected the NUDCD1 protein (arrow).

NUDCD1 Antibody (Center) - Background

NUDCD1 (CML66) contains 1 CS domain. It may play an oncogenic role in ways of favoring tumor cells proliferation, invasion and metastasis-associated with multiple pathways. NUDCD1 is expressed in leukemias and a variety of solid tumor cell lines. Northern blot analysis showed expression in normal tissues restricted to testis and heart, and no expression was found in hematopoietic tissues.

NUDCD1 Antibody (Center) - References

Wang, Q., et al. Cancer Lett. 269(1):127-138(2008) Jin, S., et al. Cancer Sci. 99(8):1670-1678(2008) Yan, Y., et al. J. Immunol. 172(1):651-660(2004) Yang, X.F., et al. Proc. Natl. Acad. Sci. U.S.A. 98(13):7492-7497(2001)