10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Mouse Kdr Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14955b

Specification

Mouse Kdr Antibody (C-term) - Product Information

Application	**WB,E**
Primary Accession	P35918
Reactivity	**Mouse**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**150462**
Antigen Region	**920-949**

Mouse Kdr Antibody (C-term) - Additional Information

Gene ID 16542

Other Names
Vascular endothelial growth factor receptor 2, VEGFR-2, Fetal liver kinase 1, FLK-1, Kinase NYK, Protein-tyrosine kinase receptor flk-1, CD309, Kdr, Flk-1, Flk1

Target/Specificity
This Mouse Kdr antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 920-949 amino acids from the C-terminal region of mouse Kdr.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
Mouse Kdr Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Kdr Antibody (C-term) - Protein Information

Name Kdr {ECO:0000312|MGI:MGI:96683}

Synonyms Flk-1, Flk1

Function Tyrosine-protein kinase that acts as a cell-surface receptor for VEGFA, VEGFC and VEGFD. Plays an essential role in the regulation of angiogenesis, vascular development, vascular permeability, and embryonic hematopoiesis. Promotes proliferation, survival, migration and differentiation of endothelial cells. Promotes reorganization of the actin cytoskeleton. Isoforms lacking a transmembrane domain, such as isoform 2, may function as decoy receptors for VEGFA, VEGFC and/or VEGFD. Isoform 2 plays an important role as a negative regulator of VEGFA- and VEGFC-mediated lymphangiogenesis by limiting the amount of free VEGFA and/or VEGFC and by preventing their binding to FLT4. Modulates FLT1 and FLT4 signaling by forming heterodimers. Binding of vascular growth factors to isoform 1 leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5- trisphosphate and the activation of protein kinase C. Mediates activation of MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. Mediates phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, reorganization of the actin cytoskeleton and activation of PTK2/FAK1. Required for VEGFA-mediated induction of NOS2 and NOS3, leading to the production of the signaling molecule nitric oxide (NO) by endothelial cells. Phosphorylates PLCG1. Promotes phosphorylation of FYN, NCK1, NOS3, PIK3R1, PTK2/FAK1 and SRC.

Cellular Location
Cell junction. Endoplasmic reticulum {ECO:0000250|UniProtKB:P35968}. Cell membrane {ECO:0000250|UniProtKB:P35968}. Note=Colocalizes with ERN1 and XBP1 in the endoplasmic reticulum in endothelial cells in a vascular endothelial growth factor (VEGF)-dependent manner (By similarity) Localized with RAP1A at cell-cell junctions {ECO:0000250|UniProtKB:P35968} [Isoform 2]: Secreted.

Tissue Location
Expressed in endothelial cells (at protein level). Detected in embryonic endothelial cells, as well as hematopoietic stem and progenitor cells. Detected in vascular endothelium. Expressed at high levels in adult heart, lung, kidney, brain and skeletal muscle, but is also expressed at lower levels in most other adult tissues


Mouse Kdr Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Mouse Kdr Antibody (C-term) - Images

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Mouse Kdr Antibody (C-term) (Cat. #AP14955b) western blot analysis in mouse lung tissue lysates (35ug/lane).This demonstrates the Kdr antibody detected the Kdr protein (arrow).

Mouse Kdr Antibody (C-term) - Background

Receptor for VEGF or VEGFC. Has a tyrosine-protein kinase activity. The VEGF-kinase ligand/receptor signaling system plays a key role in vascular development and regulation of vascular permeability.