10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Mouse Map4k2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14965C

Specification

Mouse Map4k2 Antibody (Center) - Product Information

Application	**WB,E**
Primary Accession	Q61161
Other Accession	NP_033032.1
Reactivity	**Mouse**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**91265**
Antigen Region	**432-459**

Mouse Map4k2 Antibody (Center) - Additional Information

Gene ID 26412

Other Names
Mitogen-activated protein kinase kinase kinase kinase 2, Germinal center kinase, GCK, MAPK/ERK kinase kinase kinase 2, MEK kinase kinase 2, MEKKK 2, Rab8-interacting protein, Map4k2, Rab8ip

Target/Specificity
This Mouse Map4k2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 432-459 amino acids from the Central region of mouse Map4k2.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
Mouse Map4k2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Map4k2 Antibody (Center) - Protein Information

Name Map4k2

Synonyms Rab8ip

Function Serine/threonine-protein kinase which acts as an essential component of the MAP kinase signal transduction pathway (PubMed:8643544). Acts as a MAPK kinase kinase kinase (MAP4K) and is an upstream activator of the stress-activated protein kinase/c-Jun N- terminal kinase (SAP/JNK) signaling pathway and to a lesser extent of the p38 MAPKs signaling pathway (By similarity). Required for the efficient activation of JNKs by TRAF6-dependent stimuli, including pathogen-associated molecular patterns (PAMPs) such as polyinosine- polycytidine (poly(IC)), lipopolysaccharides (LPS), lipid A, peptidoglycan (PGN), or bacterial flagellin (By similarity). To a lesser degree, IL-1 and engagement of CD40 also stimulate MAP4K2- mediated JNKs activation (By similarity). The requirement for MAP4K2/GCK is most pronounced for LPS signaling, and extends to LPS stimulation of c-Jun phosphorylation and induction of IL-8 (By similarity). Enhances MAP3K1 oligomerization, which may relieve N- terminal mediated MAP3K1 autoinhibition and lead to activation following autophosphorylation (By similarity). Mediates also the SAP/JNK signaling pathway and the p38 MAPKs signaling pathway through activation of the MAP3Ks MAP3K10/MLK2 and MAP3K11/MLK3 (By similarity). May play a role in the regulation of vesicle targeting or fusion (By similarity).


Cellular Location
Cytoplasm. Basolateral cell membrane; Peripheral membrane protein. Golgi apparatus membrane; Peripheral membrane protein

Mouse Map4k2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Mouse Map4k2 Antibody (Center) - Images



Mouse Map4k2 Antibody (Center) (Cat. #AP14965c) western blot analysis in mouse heart tissue lysates (35ug/lane).This demonstrates the Map4k2 antibody detected the Map4k2 protein (arrow).

Mouse Map4k2 Antibody (Center) - Background

Enhances MAP3K1 oligomerization, which may relieve amino-terminal mediated MAP3K1 autoinhibition and lead to activation following autophosphorylation. May play a role in the regulation of vesicle targeting or fusion.

Mouse Map4k2 Antibody (Center) - References

Zhong, J., et al. Proc. Natl. Acad. Sci. U.S.A. 106(11):4372-4377(2009)
Ren, M., et al. Proc. Natl. Acad. Sci. U.S.A. 93(10):5151-5155(1996)