10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MXI1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14974c

Specification

MXI1 Antibody (Center) - Product Information

Application	**WB,E**
Primary Accession	P50539
Other Accession	O09015, P50540, NP_005953.4, NP_569157.2, NP_001008541.1
Reactivity	**Human, Mouse**
Predicted	**Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**26062**
Antigen Region	**53-82**

MXI1 Antibody (Center) - Additional Information

Gene ID 4601

Other Names
Max-interacting protein 1, Max interactor 1, Class C basic helix-loop-helix protein 11, bHLHc11, MXI1, BHLHC11

Target/Specificity
This MXI1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 53-82 amino acids from the Central region of human MXI1.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
MXI1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

MXI1 Antibody (Center) - Protein Information

Name MXI1

Synonyms BHLHC11

Function Transcriptional repressor. MXI1 binds with MAX to form a sequence-specific DNA-binding protein complex which recognizes the core sequence 5'-CAC[GA]TG-3'. MXI1 thus antagonizes MYC transcriptional activity by competing for MAX.

Cellular Location
Nucleus.


Tissue Location
High levels found in the brain, heart and lung while lower levels are seen in the liver, kidney and skeletal muscle

MXI1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MXI1 Antibody (Center) - Images



Anti-MXI1 Antibody (Center) at 1:2000 dilution + human lung lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 26 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



MXI1 Antibody (Center) (Cat. #AP14974c) western blot analysis in mouse heart tissue lysates (35ug/lane).This demonstrates the MXI1 antibody detected the MXI1 protein (arrow).

MXI1 Antibody (Center) - Background

Expression of the c-myc gene, which produces an oncogenic transcription factor, is tightly regulated in normal cells but is frequently deregulated in human cancers. The protein encoded by this gene is a transcriptional repressor thought to negatively regulate MYC function, and is therefore a potential tumor suppressor. This protein inhibits the transcriptional activity of MYC by competing for MAX, another basic helix-loop-helix protein that binds to MYC and is required for its function. Defects in this gene are frequently found in patients with prostate tumors. Three alternatively spliced transcripts encoding different isoforms have been described. Additional alternatively spliced transcripts may exist but the products of these transcripts have not been verified experimentally.

MXI1 Antibody (Center) - References

Lofstedt, T., et al. Exp. Cell Res. 315(11):1924-1936(2009)
Baranzini, S.E., et al. Hum. Mol. Genet. 18(4):767-778(2009)
Tsao, C.C., et al. Cancer Biol. Ther. 7(10):1619-1627(2008)
Suo, X.H., et al. Zhonghua Xue Ye Xue Za Zhi 28(11):745-749(2007)
Dugast-Darzacq, C., et al. FEBS J. 274(17):4643-4653(2007)