

**CYGB Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP16031a**

**Specification**

---

**CYGB Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q8WWM9</a>
Other Accession	<a href="#">NP_599030.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	21405
Antigen Region	1-30

**CYGB Antibody (N-term) - Additional Information**

**Gene ID** 114757

**Other Names**

Cytoglobin, Histoglobin, HGb, Stellate cell activation-associated protein, CYGB, STAP

**Target/Specificity**

This CYGB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human CYGB.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CYGB Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CYGB Antibody (N-term) - Protein Information**

**Name** CYGB ([HGNC:16505](#))

**Function** Probable multifunctional globin with a hexacoordinated heme iron required for the catalysis of various reactions depending on redox condition of the cell as well as oxygen

availability (PubMed:[12359339](#), PubMed:[11893755](#), PubMed:[19147491](#), PubMed:[20511233](#), PubMed:[28671819](#), PubMed:[28393874](#), PubMed:[29128400](#), PubMed:[33576020](#), PubMed:[34930834](#), PubMed:[15165856](#)). Has a nitric oxide dioxygenase (NOD) activity and is most probably involved in cell-mediated and oxygen-dependent nitric oxide consumption (PubMed:[19147491](#), PubMed:[20511233](#), PubMed:[28671819](#), PubMed:[28393874](#)). By scavenging this second messenger may regulate several biological processes including endothelium-mediated vasodilation and vascular tone (PubMed:[19147491](#), PubMed:[28393874](#)). Under normoxic conditions functions as a nitric oxide dioxygenase (NOD) but under hypoxic conditions the globin may switch its function to that of a nitrite (NO<sub>2</sub>) reductase (NiR), generating nitric oxide (PubMed:[29128400](#)). Could also have peroxidase and superoxide dismutase activities, detoxifying reactive oxygen species and protecting cells against oxidative stress (PubMed:[12359339](#), PubMed:[33576020](#), PubMed:[34930834](#)). Also binds dioxygen with low affinity and could function as an oxygen sensor but has probably no function as a respiratory oxygen carrier (PubMed:[11893755](#), PubMed:[15299006](#), PubMed:[20553503](#)).

#### Cellular Location

Cytoplasm. Nucleus

#### Tissue Location


Widely expressed. Highest expression in heart, stomach, bladder and small intestine.

#### CYGB Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### CYGB Antibody (N-term) - Images



CYGB Antibody (N-term) (Cat. #AP16031a) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the CYGB antibody detected the CYGB protein (arrow).

#### CYGB Antibody (N-term) - Background

Cytoglobin is a ubiquitously expressed hexacoordinate hemoglobin that may facilitate diffusion of oxygen through tissues, scavenge nitric oxide or other reactive oxygen species, or serve a protective function during oxidative stress (Trent and Hargrove, 2002 [PubMed 11893755]).

#### **CYGB Antibody (N-term) - References**

Gardner, A.M., et al. J. Biol. Chem. 285(31):23850-23857(2010)  
Lechauve, C., et al. FEBS J. 277(12):2696-2704(2010)  
Shaw, R.J., et al. Br. J. Cancer 101(1):139-144(2009)  
Halligan, K.E., et al. J. Biol. Chem. 284(13):8539-8547(2009)  
Ostojic, J., et al. Arch. Ophthalmol. 126(11):1530-1536(2008)