

**ZIC1 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP16034a****Specification**

---

**ZIC1 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q15915</a>
Other Accession	<a href="#">P46684</a> , <a href="#">NP_003403.2</a>
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	48309
Antigen Region	6-34

**ZIC1 Antibody (N-term) - Additional Information****Gene ID** 7545**Other Names**

Zinc finger protein ZIC 1, Zinc finger protein 201, Zinc finger protein of the cerebellum 1, ZIC1, ZIC, ZNF201

**Target/Specificity**

This ZIC1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 6-34 amino acids from the N-terminal region of human ZIC1.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ZIC1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**ZIC1 Antibody (N-term) - Protein Information****Name** ZIC1

### Synonyms ZIC, ZNF201

**Function** Acts as a transcriptional activator. Involved in neurogenesis. Plays important roles in the early stage of organogenesis of the CNS, as well as during dorsal spinal cord development and maturation of the cerebellum. Involved in the spatial distribution of mossy fiber (MF) neurons within the pontine gray nucleus (PGN). Plays a role in the regulation of MF axon pathway choice. Promotes MF migration towards ipsilaterally-located cerebellar territories. May have a role in shear flow mechanotransduction in osteocytes. Retains nuclear GLI1 and GLI3 in the cytoplasm. Binds to the minimal GLI-consensus sequence 5'-TGGGTGGTC-3' (By similarity).

### Cellular Location

Nucleus. Cytoplasm. Note=Localizes in the cytoplasm in presence of MDFIC overexpression.

### Tissue Location


CNS. A high level expression is seen in the cerebellum. Detected in the nuclei of the cerebellar granule cell lineage from the progenitor cells of the external germinal layer to the postmigrated cells of the internal granular layer. Detected in medulloblastoma (26/29 cases), but not present in all other tumors examined.

### ZIC1 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### ZIC1 Antibody (N-term) - Images



ZIC1 Antibody (N-term) (Cat. #AP16034a) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the ZIC1 antibody detected the ZIC1 protein (arrow).



All lanes : Anti-ZIC1 Antibody (N-term) at 1:2000 dilution Lane 1: Jurkat whole cell lysates Lane 2: SH-SY5Y whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 48 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

#### **ZIC1 Antibody (N-term) - Background**

This gene encodes a member of the ZIC family of C2H2-type zinc finger proteins. Members of this family are important during development. Aberrant expression of this gene is seen in medulloblastoma, a childhood brain tumor. This gene is closely linked to the gene encoding zinc finger protein of the cerebellum 4, a related family member on chromosome 3. This gene encodes a transcription factor that can bind and transactivate the apolipoprotein E gene.

#### **ZIC1 Antibody (N-term) - References**

- Brill, E., et al. Cancer Res. 70(17):6891-6901(2010)
- Wang, L.J., et al. Biochem. Biophys. Res. Commun. 379(4):959-963(2009)
- Baranzini, S.E., et al. Hum. Mol. Genet. 18(4):767-778(2009)
- Arora, A., et al. RNA 14(7):1290-1296(2008)
- Pourebrahim, R., et al. FEBS Lett. 581(26):5122-5126(2007)