10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Mouse Cdk4 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16073b

Specification

Mouse Cdk4 Antibody (C-term) - Product Information

Application	**WB,E**
Primary Accession	P30285
Other Accession	P35426, NP_034000.1
Reactivity	**Mouse**
Predicted	**Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**33751**
Antigen Region	**219-246**

Mouse Cdk4 Antibody (C-term) - Additional Information

Gene ID 12567

Other Names
Cyclin-dependent kinase 4, CRK3, Cell division protein kinase 4, PSK-J3, Cdk4, Crk3

Target/Specificity
This Mouse Cdk4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 219-246 amino acids from the C-terminal region of mouse Cdk4.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
Mouse Cdk4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Cdk4 Antibody (C-term) - Protein Information

Name Cdk4

Synonyms Crk3

Function Ser/Thr-kinase component of cyclin D-CDK4 (DC) complexes that phosphorylate and inhibit members of the retinoblastoma (RB) protein family including RB1 and regulate the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complexes and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenenic and antimitogenic signals. Also phosphorylates SMAD3 in a cell-cycle-dependent manner and represses its transcriptional activity. Component of the ternary complex, cyclin D/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (By similarity).


Cellular Location
Cytoplasm {ECO:0000250|UniProtKB:P11802}. Nucleus {ECO:0000250|UniProtKB:P11802}. Nucleus membrane {ECO:0000250|UniProtKB:P11802}. Note=Cytoplasmic when non-complexed Forms a cyclin D-CDK4 complex in the cytoplasm as cells progress through G(1) phase. The complex accumulates on the nuclear membrane and enters the nucleus on transition from G(1) to S phase. Also present in nucleoli and heterochromatin lumps. Colocalizes with RB1 after release into the nucleus (By similarity). {ECO:0000250|UniProtKB:P11802}

Mouse Cdk4 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Mouse Cdk4 Antibody (C-term) - Images



Mouse Cdk4 Antibody (C-term) (Cat. #AP16073b) western blot analysis in mouse NIH-3T3 cell line lysates (35ug/lane).This demonstrates the Cdk4 antibody detected the Cdk4 protein (arrow).

Mouse Cdk4 Antibody (C-term) - Background

Cdk4 is probably involved in the control of the cell cycle.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Mouse Cdk4 Antibody (C-term) - References

Trivedi, C.M., et al. Dev. Cell 19(3):450-459(2010)
Beilke, S., et al. Oncogene 29(28):4058-4067(2010)
Puyol, M., et al. Cancer Cell 18(1):63-73(2010)
Wiedemeyer, W.R., et al. Proc. Natl. Acad. Sci. U.S.A. 107(25):11501-11506(2010)
Michaud, K., et al. Cancer Res. 70(8):3228-3238(2010)