

NCOA4 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16218b**Specification**

NCOA4 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q13772
Other Accession	NP_005428.1 , NP_001138732.1 , NP_001138734.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	69726
Antigen Region	496-524

NCOA4 Antibody (C-term) - Additional Information**Gene ID** 8031**Other Names**

Nuclear receptor coactivator 4, NCoA-4, Androgen receptor coactivator 70 kDa protein, 70 kDa AR-activator, 70 kDa androgen receptor coactivator, Androgen receptor-associated protein of 70 kDa, Ret-activating protein ELE1, NCOA4, ARA70, ELE1, RFG

Target/Specificity

This NCOA4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 496-524 amino acids from the C-terminal region of human NCOA4.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NCOA4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NCOA4 Antibody (C-term) - Protein Information**Name** NCOA4

Synonyms ARA70, ELE1, RFG

Function Enhances the androgen receptor transcriptional activity in prostate cancer cells. Ligand-independent coactivator of the peroxisome proliferator-activated receptor (PPAR) gamma.

Tissue Location


Widely expressed. Also detected in adipose tissues and in different cell lines. Isoform Beta is only expressed in testis

NCOA4 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NCOA4 Antibody (C-term) - Images



NCOA4 Antibody (C-term) (Cat. #AP16218b) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the NCOA4 antibody detected the NCOA4 protein (arrow).

NCOA4 Antibody (C-term) - Background

This gene encodes an androgen receptor coactivator. The encoded protein interacts with the androgen receptor in a ligand-dependent manner to enhance its transcriptional activity. Chromosomal translocations between this gene and the ret tyrosine kinase gene, also located on chromosome 10, have been associated with papillary thyroid carcinoma.

NCOA4 Antibody (C-term) - References

Sadow, P.M., et al. Endocr. Pathol. 21(2):73-79(2010)
Landa, I., et al. PLoS Genet. 5 (9), E1000637 (2009) :

Richardson, D.S., et al. Cancer Res. 69(11):4861-4869(2009)
Peng, Y., et al. Am. J. Pathol. 172(1):225-235(2008)
Olsen, J.V., et al. Cell 127(3):635-648(2006)