

FAM38B Antibody-(C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16313b**Specification**

FAM38B Antibody-(C-term) - Product Information

Application	WB,E
Primary Accession	Q9H5I5
Other Accession	Q8CD54 , NP_071351.2
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	2575-2604

FAM38B Antibody-(C-term) - Additional Information**Gene ID** 63895**Other Names**

Piezo-type mechanosensitive ion channel component 2, Protein FAM38B, PIEZO2, C18orf30, C18orf58, FAM38B

Target/Specificity

This FAM38B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 2575-2604 amino acids of human FAM38B.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FAM38B Antibody-(C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FAM38B Antibody-(C-term) - Protein Information**Name** PIEZO2 ([HGNC:26270](#))**Function** Pore-forming subunit of a mechanosensitive non-specific cation channel required for

rapidly adapting mechanically activated (MA) currents and has a key role in sensing touch and tactile pain. Required for Merkel-cell mechanotransduction. Plays a major role in light-touch mechanosensation.

Cellular Location


Cell membrane {ECO:0000250|UniProtKB:Q8CD54}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q8CD54}

FAM38B Antibody-(C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FAM38B Antibody-(C-term) - Images



All lanes : Anti-FAM38B Antibody-(C-term) at 1:500 dilution Lane 1: HepG2 whole cell lysate Lane 2: A549 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Observed band size : 318kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-FAM38B Antibody-(C-term) at 1:500 dilution Lane 1: HeLa whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: A549 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Observed band size : 318kDa Blocking/Dilution buffer: 5% NFDm/TBST.

FAM38B Antibody-(C-term) - Background

Piezos are large transmembrane proteins conserved among various species, all having between 24 and 36 predicted transmembrane domains. 'Piezo' comes from the Greek 'piesi,' meaning 'pressure.' The PIEZO2 protein has a role in rapidly adapting mechanically activated (MA) currents in somatosensory neurons (Coste et al., 2010 [PubMed 20813920]).

FAM38B Antibody-(C-term) - References

Coste, B., et al. Science 330(6000):55-60(2010)
Brandenberger, R., et al. Nat. Biotechnol. 22(6):707-716(2004)

FAM38B Antibody-(C-term) - Citations

- [Linoleic acid improves PIEZO2 dysfunction in a mouse model of Angelman Syndrome](#)
- [The mechanosensitive Piezo1 channel mediates heart mechano-chemo transduction](#)