

RNF34 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16633b**Specification**

RNF34 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q969K3
Other Accession	Q6AYH3 , Q99KR6 , Q5E9J6 , NP_079402.2 , NP_919247.1
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	41641
Antigen Region	283-311

RNF34 Antibody (C-term) - Additional Information**Gene ID** 80196**Other Names**

E3 ubiquitin-protein ligase RNF34, 632- {ECO:0000269|PubMed:25012219, ECO:0000269|Ref13}, Caspase regulator CARP1, Caspases-8 and -10-associated RING finger protein 1, CARP-1, FYVE-RING finger protein Momo, RNF34 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=17297)
HGNC:17297

Target/Specificity

This RNF34 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 283-311 amino acids from the C-terminal region of human RNF34.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RNF34 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RNF34 Antibody (C-term) - Protein Information

Name RNF34 ([HGNC:17297](#))

Function E3 ubiquitin-protein ligase that regulates several biological processes through the ubiquitin-mediated proteasomal degradation of various target proteins. Ubiquitinates the caspases CASP8 and CASP10, promoting their proteasomal degradation, to negatively regulate cell death downstream of death domain receptors in the extrinsic pathway of apoptosis (PubMed:[15069192](#)). May mediate 'Lys-48'-linked polyubiquitination of RIPK1 and its subsequent proteasomal degradation thereby indirectly regulating the tumor necrosis factor-mediated signaling pathway (Ref.13). Negatively regulates p53/TP53 through its direct ubiquitination and targeting to proteasomal degradation (PubMed:[17121812](#)). Indirectly, may also negatively regulate p53/TP53 through ubiquitination and degradation of SFN (PubMed:[18382127](#)). Mediates PPARGC1A proteasomal degradation probably through ubiquitination thereby indirectly regulating the metabolism of brown fat cells (PubMed:[22064484](#)). Possibly involved in innate immunity, through 'Lys-48'-linked polyubiquitination of NOD1 and its subsequent proteasomal degradation (PubMed:[25012219](#)).

Cellular Location

Cell membrane; Peripheral membrane protein. Endomembrane system {ECO:0000250|UniProtKB:Q6AYH3}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q6AYH3}. Nucleus Nucleus speckle. Cytoplasm, cytosol


Tissue Location

Ubiquitous. Detected in heart, brain, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, colon and leukocytes.

RNF34 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RNF34 Antibody (C-term) - Images

RNF34 Antibody (C-term) (Cat. #AP16633b) western blot analysis in Hela cell line lysates (35ug/lane). This demonstrates the RNF34 antibody detected the RNF34 protein (arrow).

RNF34 Antibody (C-term) - Background

The protein encoded by this gene contains a RINF finger, a motif known to be involved in protein-protein and protein-DNA interactions. This protein interacts with DNAJA3/hTid-1, which is a DnaJ protein reported to function as a modulator of apoptosis. Overexpression of this gene in Hela cells was shown to confer the resistance to TNF-alpha induced apoptosis, suggesting an anti-apoptotic function of this protein. This protein can be cleaved by caspase-3 during the induction of apoptosis. Alternatively spliced transcript variants encoding distinct isoforms have been reported.

RNF34 Antibody (C-term) - References

Erlbruch, A., et al. Proteomics 10(16):2890-2900(2010)
Yang, W., et al. Cell Cycle 7(5):670-682(2008)
Yang, W., et al. J. Biol. Chem. 282(5):3273-3281(2007)
Konishi, T., et al. Mol. Cancer Ther. 4(5):743-750(2005)
Sasaki, S., et al. J. Exp. Clin. Cancer Res. 23(3):507-512(2004)