

SULT2A1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16646a

Specification

SULT2A1 Antibody (N-term) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region WB,E <u>Q06520</u> <u>P52842</u>, <u>NP_003158.2</u> Human Monkey Rabbit Polyclonal Rabbit IgG 33780 2-30

SULT2A1 Antibody (N-term) - Additional Information

Gene ID 6822

Other Names

Bile salt sulfotransferase, Dehydroepiandrosterone sulfotransferase, DHEA-ST, Hydroxysteroid Sulfotransferase, HST, ST2, ST2A3, Sulfotransferase 2A1, ST2A1, SULT2A1, HST, STD

Target/Specificity

This SULT2A1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 2-30 amino acids from the N-terminal region of human SULT2A1.

Dilution WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SULT2A1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SULT2A1 Antibody (N-term) - Protein Information

Name SULT2A1



Synonyms HST, STD

Function Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the sulfonation of steroids and bile acids in the liver and adrenal glands. Mediates the sulfation of a wide range of steroids and sterols, including pregnenolone, androsterone, DHEA, bile acids, cholesterol and as well many xenobiotics that contain alcohol and phenol functional groups (PubMed:<u>7678732</u>, PubMed:<u>2268288</u>, PubMed:<u>14573603</u>, PubMed:<u>18042734</u>, PubMed:<u>19589875</u>, PubMed:<u>21187059</u>, PubMed:<u>29671343</u>, PubMed:<u>7854148</u>). Sulfonation increases the water solubility of most compounds, and therefore their renal excretion, but it can also result in bioactivation to form active metabolites. Plays an important role in maintening steroid and lipid homeostasis (PubMed:<u>21187059</u>, PubMed:<u>19589875</u>, PubMed:<u>14573603</u>). Plays a key role in bile acid metabolism (PubMed:<u>2268288</u>). In addition, catalyzes the metabolic activation of potent carcinogenic polycyclic arylmethanols (By similarity).

Cellular Location Cytoplasm.

Tissue Location

Liver, adrenal and at lower level in the kidney. Is present in human fetus in higher level in the adrenal than the liver and the kidney

SULT2A1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

NCI-H292 95 72 55 36 28

SULT2A1 Antibody (N-term) (Cat. #AP16646a) western blot analysis in NCI-H292 cell line lysates (35ug/lane).This demonstrates the SULT2A1 antibody detected the SULT2A1 protein (arrow).

SULT2A1 Antibody (N-term) - Background

SULT2A1 Antibody (N-term) - Images



This gene encodes a member of the sulfotransferase family. Sulfotransferases aid in the metabolism of drugs and endogenous compounds by converting these substances into more hydrophilic water-soluble sulfate conjugates that can be easily excreted. This protein catalyzes the sulfation of steroids and bile acids in the liver and adrenal glands, and may have a role in the inherited adrenal androgen excess in women with polycystic ovary syndrome.

SULT2A1 Antibody (N-term) - References

Huang, J., et al. Xenobiotica 40(3):184-194(2010) Li, J., et al. Breast Cancer Res. 12 (2), R19 (2010) : Senggunprai, L., et al. Drug Metab. Dispos. 37(8):1711-1717(2009) Chakrabarti, B., et al. Autism Res 2(3):157-177(2009) Saito, A., et al. J. Hum. Genet. 54(6):317-323(2009)