

OR10S1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16691b

Specification

OR10S1 Antibody (C-term) - Product Information

Application WB,E
Primary Accession Q8NGN2

Other Accession NP_001004474.1

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region
Reactivity
Human
Rabbit
Polyclonal
Rabbit IgG
226-254

OR10S1 Antibody (C-term) - Additional Information

Gene ID 219873

Other Names

Olfactory receptor 10S1, Olfactory receptor OR11-279, OR10S1

Target/Specificity

This OR10S1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 226-254 amino acids from the C-terminal region of human OR10S1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR10S1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR10S1 Antibody (C-term) - Protein Information

Name OR10S1

Function Odorant receptor.



Cellular Location


Cell membrane; Multi-pass membrane protein.

OR10S1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

OR10S1 Antibody (C-term) - Images



OR10S1 Antibody (C-term) (Cat. #AP16691b) western blot analysis in Hela cell line lysates (35ug/lane). This demonstrates the OR10S1 antibody detected the OR10S1 protein (arrow).

OR10S1 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR10S1 Antibody (C-term) - References





Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004) Fuchs, T., et al. Genomics 80(3):295-302(2002)