

#### **TGFBI Antibody (N-term)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16708a

### **Specification**

### TGFBI Antibody (N-term) - Product Information

Application WB, FC,E
Primary Accession 015582

Other Accession <u>Q95215</u>, <u>Q11780</u>, <u>P82198</u>, <u>NP 000349.1</u>,

P55906

Reactivity Human

Predicted Bovine, Mouse, Pig, Rabbit

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 74681
Antigen Region 106-135

# TGFBI Antibody (N-term) - Additional Information

# **Gene ID** 7045

#### **Other Names**

Transforming growth factor-beta-induced protein ig-h3, Beta ig-h3, Kerato-epithelin, RGD-containing collagen-associated protein, RGD-CAP, TGFBI, BIGH3

### Target/Specificity

This TGFBI antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 106-135 amino acids from the N-terminal region of human TGFBI.

### **Dilution**

WB~~1:1000 FC~~1:10~50

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at  $2-8^{\circ}$ C for up to 2 weeks. For long term storage store at  $-20^{\circ}$ C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

TGFBI Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### TGFBI Antibody (N-term) - Protein Information



## **Name TGFBI**

#### **Synonyms** BIGH3

**Function** Plays a role in cell adhesion (PubMed:<u>8024701</u>). May play a role in cell-collagen interactions (By similarity).

#### **Cellular Location**

Secreted. Secreted, extracellular space, extracellular matrix Note=May be associated both with microfibrils and with the cell surface (PubMed:8077289).

### **Tissue Location**


Highly expressed in the corneal epithelium (PubMed:27609313, PubMed:8077289). Expressed in heart, placenta, lung, liver, skeletal muscle, kidney and pancreas (PubMed:8077289)

### TGFBI Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

### TGFBI Antibody (N-term) - Images



Western blot analysis of TGFBI (arrow) using rabbit polyclonal TGFBI Antibody (N-term) (Cat. #AP16708a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the TGFBI gene.





TGFBI Antibody (N-term) (Cat. #AP16708a) flow cytometric analysis of U251 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

# TGFBI Antibody (N-term) - Background

This gene encodes an RGD-containing protein that binds to type I, II and IV collagens. The RGD motif is found in many extracellular matrix proteins modulating cell adhesion and serves as a ligand recognition sequence for several integrins. This protein plays a role in cell-collagen interactions and may be involved in endochondrial bone formation in cartilage. The protein is induced by transforming growth factor-beta and acts to inhibit cell adhesion. Mutations in this gene are associated with multiple types of corneal dystrophy.

### TGFBI Antibody (N-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010) Edelstein, S.L., et al. Cornea 29(6):698-700(2010) Romero, P., et al. Mol. Vis. 16, 1601-1609 (2010): Paliwal, P., et al. Mol. Vis. 16, 1429-1438 (2010): Yang, J., et al. Mol. Vis. 16, 1186-1193 (2010):