

IL28B Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16853a

Specification

IL28B Antibody (N-term) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW Antigen Region WB,E <u>O8IZI9</u> <u>O8IZJ0</u>, <u>NP_742151.2</u> Human Rabbit Polyclonal Rabbit IgG 21706 34-63

IL28B Antibody (N-term) - Additional Information

Gene ID 282617

Other Names

Interferon lambda-3, IFN-lambda-3, Cytokine Zcyto22, Interleukin-28B, IL-28B, Interleukin-28C, IL-28C, IFNL3, IL28B, IL28C, ZCYTO22

Target/Specificity

This IL28B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 34-63 amino acids from the N-terminal region of human IL28B.

Dilution WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IL28B Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

IL28B Antibody (N-term) - Protein Information

Name IFNL3

Synonyms IL28B, IL28C, ZCYTO22



Function Cytokine with antiviral, antitumour and immunomodulatory activities. Plays a critical role in the antiviral host defense, predominantly in the epithelial tissues. Acts as a ligand for the heterodimeric class II cytokine receptor composed of IL10RB and IFNLR1, and receptor engagement leads to the activation of the JAK/STAT signaling pathway resulting in the expression of IFN-stimulated genes (ISG), which mediate the antiviral state. Has a restricted receptor distribution and therefore restricted targets: is primarily active in epithelial cells and this cell type-selective action is because of the epithelial cell-specific expression of its receptor IFNLR1. Seems not to be essential for early virus-activated host defense in vaginal infection, but plays an important role in Toll-like receptor (TLR)- induced antiviral defense. Plays a significant role in the antiviral immune defense in the intestinal epithelium. Exerts an immunomodulatory effect by up-regulating MHC class I antigen expression.

Cellular Location Secreted.

IL28B Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

IL28B Antibody (N-term) - Images



IL28B Antibody (N-term) (Cat. #AP16853a) western blot analysis in SK-BR-3 cell line lysates (35ug/lane).This demonstrates the IL28B antibody detected the IL28B protein (arrow).





Western blot analysis of IL28B (arrow) using rabbit polyclonal IL28B Antibody (N-term) (Cat. #AP16853a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the IL28B gene.

IL28B Antibody (N-term) - Background

This gene encodes a cytokine distantly related to type I interferons and the IL-10 family. This gene, interleukin 28A (IL28A), and interleukin 29 (IL29) are three closely related cytokine genes that form a cytokine gene cluster on a chromosomal region mapped to 19q13. Expression of the cytokines encoded by the three genes can be induced by viral infection. All three cytokines have been shown to interact with a heterodimeric class II cytokine receptor that consists of interleukin 10 receptor, beta (IL10RB) and interleukin 28 receptor, alpha (IL28RA).

IL28B Antibody (N-term) - References

Lindh, M., et al. Hepatology 52(5):1860-1861(2010) Pineda, J.A., et al. Clin. Infect. Dis. 51(7):788-795(2010) Grebely, J., et al. Hepatology 52(4):1216-1224(2010) Stattermayer, A.F., et al. Clin. Gastroenterol. Hepatol. (2010) In press : Iwata, R., et al. J. Viral Hepat. (2010) In press : **IL28B Antibody (N-term) - Citations**

• <u>Cellular Mechanism for Impaired Hepatitis C Virus Clearance by Interferon Associated with</u> <u>IFNL3 Gene Polymorphisms Relates to Intrahepatic Interferon-λ Expression.</u>