10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MAP3K7IP3 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP17071a

Specification

MAP3K7IP3 Antibody (N-term) - Product Information

Application	**WB,E**
Primary Accession	Q8N5C8
Other Accession	Q571K4, NP_690000.2
Reactivity	**Human**
Predicted	**Mouse**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**78653**
Antigen Region	**1-30**

MAP3K7IP3 Antibody (N-term) - Additional Information

Gene ID 257397

Other Names
TGF-beta-activated kinase 1 and MAP3K7-binding protein 3, Mitogen-activated protein kinase kinase kinase 7-interacting protein 3, NF-kappa-B-activating protein 1, TAK1-binding protein 3, TAB-3, TGF-beta-activated kinase 1-binding protein 3, TAB3, MAP3K7IP3

Target/Specificity
This MAP3K7IP3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human MAP3K7IP3.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
MAP3K7IP3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MAP3K7IP3 Antibody (N-term) - Protein Information

Name TAB3 {ECO:0000303|PubMed:14633987, ECO:0000312|HGNC:HGNC:30681}

Function Adapter required to activate the JNK and NF-kappa-B signaling pathways through the specific recognition of 'Lys-63'-linked polyubiquitin chains by its RanBP2-type zinc finger (NZF) (PubMed:14633987, PubMed:14766965, PubMed:15327770, PubMed:22158122). Acts as an adapter linking MAP3K7/TAK1 and TRAF6 to 'Lys-63'-linked polyubiquitin chains (PubMed:14633987, PubMed:14766965, PubMed:15327770, PubMed:22158122, PubMed:36593296). The RanBP2-type zinc finger (NZF) specifically recognizes Lys-63'-linked polyubiquitin chains unanchored or anchored to the substrate proteins such as RIPK1/RIP1 and RIPK2: this acts as a scaffold to organize a large signaling complex to promote autophosphorylation of MAP3K7/TAK1, and subsequent activation of I-kappa-B-kinase (IKK) core complex by MAP3K7/TAK1 (PubMed:15327770, PubMed:18079694, PubMed:22158122).


Tissue Location
Widely expressed. Constitutively overexpressed in certain tumor tissues. [Isoform 2]: Minor transcript.

MAP3K7IP3 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MAP3K7IP3 Antibody (N-term) - Images



MAP3K7IP3 Antibody (N-term) (Cat. #AP17071a) western blot analysis in HL-60 cell line lysates (35ug/lane).This demonstrates the MAP3K7IP3 antibody detected the MAP3K7IP3 protein (arrow).

MAP3K7IP3 Antibody (N-term) - Background

The product of this gene functions in the NF-kappaB signal transduction pathway. The encoded protein, and the similar and functionally redundant protein MAP3K7IP2/TAB2, forms a ternary complex with the protein kinase MAP3K7/TAK1 and either TRAF2 or TRAF6 in response to stimulation with the pro-inflammatory

cytokines TNF or IL-1. Subsequent MAP3K7/TAK1 kinase activity triggers a signaling cascade leading to activation of the NF-kappaB transcription factor. The human genome contains a related pseudogene. Alternatively spliced transcript variants have been described, but their biological validity has not been determined.

MAP3K7IP3 Antibody (N-term) - References

Besse, A., et al. J. Biol. Chem. 282(6):3918-3928(2007)
Kanayama, A., et al. Mol. Cell 15(4):535-548(2004)
Jin, G., et al. Proc. Natl. Acad. Sci. U.S.A. 101(7):2028-2033(2004)
Cheung, P.C., et al. Biochem. J. 378 (PT 1), 27-34 (2004) :
Bouwmeester, T., et al. Nat. Cell Biol. 6(2):97-105(2004)